



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

2000/01903

REC'D 28 AUG 2000

WIPO

PCT

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

99104519.6

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

Der Präsident des Europäischen Patentamts;  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN  
THE HAGUE,

22/05/00



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung  
Sheet 2 of the certificate  
Page 2 de l'attestation**

Anmeldung Nr.: 99104519.6  
Application no.:  
Demande n°:

Anmeldetag: 06/03/99  
Date of filing:  
Date de dépôt:

Anmelder:  
Applicant(s):  
Demandeur(s):  
ARTEMIS Pharmaceuticals GmbH  
51063 Köln  
GERMANY

Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:

Stable recombinant influenza viruses free of helper viruses

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

C12N15/86, C12N7/01, C12N5/10, A61K39/00, A61K39/145, A61K48/00

Am Anmeldetag benannte Vertragstaaten:  
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:

06. März 1999

## Stable Recombinant Influenza Viruses Free of Helper Viruses

The invention relates to a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus; a method for the production of said recombinant influenza virus, a method for constructing specific ribozyme-sensitive influenza carrier strains; pharmaceutical compositions comprising said recombinant influenza viruses; and the use of said recombinant influenza virus for preparing medicaments for vaccination purposes.

Redesigning influenza virus into a vector system for expression of foreign genes similar to what has been achieved in several other thoroughly studied viruses such as adenovirus, retrovirus, Semliki Forest virus or Rabies virus has the advantage of an industrially well established mode of cheap propagation for influenza in fertilized chicken eggs leading to rather high titers (above  $10^{10}$ /ml). On the other hand none of its genes may be deleted from the influenza genome according to our present knowledge, and give room for large-size foreign insertions. Only small fragments of foreign polypeptide chains such as B cell epitopes (10 to 15 amino acids) may be inserted into selected positions within two of the viral proteins, i.e. in exchange for one of the variable antigenic regions located at the surface of hemagglutinin (Muster et al., Mucosal model of immunization against human immunodeficiency virus type 1 with a chimeric influenza virus. *J. Virol.* 69 (11), 6678-6686 (1995)), or into the stalk sequence of viral neuraminidase (Garcia-Sastre and Palese, The cytoplasmic tail of the neuraminidase protein of influenza A virus does not play an important role in the packaging of this protein into viral envelopes. *Virus Res.* 37, 37-47 (1995)), and be stably maintained as functional fusion proteins. Constructs of this kind turned out to be useful for experimental vaccination in a few cases studied, but only rather few clearly defined epitope sequences (of ten to twelve amino acids each) are known today, and some of them might also be misfolded within such restricted fusion protein positions, or in other cases interfere with the correct tertiary structure and function of their host polypeptide chains.

Incorporation of a full-size foreign protein into influenza via reverse genetics, encoded by an independent ninth vRNA molecule in addition to its regular set of eight standard vRNA segments is without special provisions only transiently possible (Luytjes et al., Amplification, expression, and packaging of a foreign gene by influenza virus. *Cell* 59, 1107-1113 (1989);

Enami et al., An influenza virus containing nine different RNA segments. *Virology* 185, 291-298 (1991)). In the absence of a continuous selective pressure any additional recombinant vRNA segment cannot be stably maintained as long as a wildtype promoter sequence is used on that ninth vRNA segment, and it will inadvertently be lost after few steps of propagation. Using a different system of influenza reverse genetics developed in our laboratory (Zobel et al., RNA polymerase I catalysed transcription of insert viral cDNA. *Nucleic Acids Res.* 21, 3607-3614 (1993); Neumann et al., RNA polymerase I-mediated expression of influenza viral RNA molecules. *Virology* 202, 477-479 (1994)), which was built around ~~in vivo synthesis of~~ recombinant vRNA molecules by cellular RNA polymerase I transcription of the respective template cDNA constructs, promoter-up mutations have been designed by nucleotide substitutions (Neumann and Hobom, Mutational analysis of influenza virus promoter elements *in vivo*. *J. Gen. Virol.* 76, 1709-1717 (1995)). When these are attached to a recombinant ninth vRNA segment its increased transcription and amplification rate will not only compensate for the losses suffered spontaneously, but even cause accumulation of the foreign vRNA segment during simple viral passaging, in the absence of any selection. However, due to its over-replication relative to all of the regular influenza vRNA segments (which of course are connected to wild-type promoter sequences) after catching up with the others the foreign segment will become over-abundant. This increasingly will result in viral particles that have incorporated several copies of recombinant vRNA, but no longer have a full set of all eight standard segments incorporated among an average of about 15 vRNA molecules present within a virion. Such particles are defective and will not result in plaque formation, hence after an initial increase of recombinant viral particles during the first steps of propagation a dramatic decrease is observed, usually at the third or fourth step of viral passaging, depending on the size of the recombinant vRNA and the level of the promoter-up mutation attached. A balanced situation with regard to the insert length and the level of promoter activity can be achieved, and has been propagated in a particular case over 11 passages, with essentially stable levels of recombinant viruses among a majority of helper viruses (around 80%) during these steps. If a full-level promoter-up mutation is used (1104 or the new variant 1920, see below) a balanced-level propagation is reached in conjunction with a recombinant vRNA size of 4000 nucleotides (unpublished).

In all of these preparations, both in transiently achieved increased yields (up to 40% of recombinants after three or four steps of viral passage), and in a balanced propagation of

recombinant influenza viruses (10 - 20%) the respective viral progeny inadvertently constitute mixtures with a majority of non-recombinant helper viruses. These result both from a statistical mode of packaging vRNA molecules into a virion (the ninth segment may not be co-packaged), and from the fraction of cells solely infected by helper virus.

The problems of fractional yields and of instability during viral propagation of recombinant influenza are the problems to be solved with the present invention.

---

~~Starting out from two observations in this laboratory which are discussed below and which~~ concern two hitherto unsuspected properties of influenza viral RNA polymerase in its interaction with terminally adapted influenza-specific RNA molecules, a new technique for the construction of stable recombinant influenza viruses was found.

As previously described in WO 96/10641 plasmid constructs are designed to generate influenza vRNA-like molecules *in vivo* by cellular RNA polymerase I transcription following plasmid DNA transfection into tissue culture cells, and to this end contain flanking rDNA promoter and terminator elements, externally located relative to any cDNA constructs in between. The resulting recombinant vRNA molecules are designed to contain 5' and 3' recognition sequences for influenza viral RNA polymerase, which however carry up to five nucleotide substitutions (in promoter-up mutant pHL1920) resulting in a substantial increase of expression over wildtype influenza promoter levels. While recombinant pseudoviral RNA is initially transcribed by RNA polymerase I, further amplification and mRNA transcription depends on the presence of viral RNA polymerase and viral nucleoprotein in the cell, which generally are provided by infection of a helper virus. As a consequence the progeny viral yield will constitute a mixture of recombinant viruses together with a majority of wild-type helper viruses.

In the new technique the recombinant vRNA-like molecules as transcribed by RNA polymerase I are constructed as ambisense RNA double segments, with one reading frame (an influenza gene) in sense and a second one (a foreign gene) in anti-sense orientation, or vice versa. In such constructs both reading frames will be expressed via the cap-snatching mode, even if at different levels. Again, infection by helper virus is required to provide the necessary viral early and late proteins for genetic expression and virion packaging. However, the particular helper virus used in the new method is a recombinant virus carrying 2x2 specifically designed ribozyme targets inserted into the flanking non-coding sequences of one

of its eight vRNA segments. The viral segment chosen to become ribozyme-sensitive is always identical to the one used in constructing the recombinant ambisense RNA molecule, as the viral carrier gene in covalent linkage with an additional foreign gene.

Recombinant influenza viruses produced in this way through RNA polymerase I transcription of an ambisense viral RNA molecule followed by infection with that specifically designed type of ribozyme-sensitive helper virus will carry one of the influenza genes twice, once within that ribozyme-sensitive helper vRNA segment, and a second time within the ribozyme-

~~resistant ambisense segment. Recombinant viruses of this type are again obtained initially~~

only as a mixture together with a majority of non-recombinant helper viruses. A progeny viral passage through tissue culture cells (293T) which before have been transiently transfected with plasmid constructs expressing the respective double-headed ribozyme will (in one step) inactivate the ribozyme-sensitive segment by a factor of up to 100. One or two rounds of such ribozyme treatment *in vivo* will at the same time (a) purify the recombinant virus from its non-recombinant helper contaminants, and (b) delete the sensitive vRNA helper segment from within the initial (additive) recombinant virus.

As a result recombinant influenza viruses are isolated along this several-step procedure, which are free of contaminating helper viruses and carry seven regular and one ambisense vRNA segments, all in a balanced replication mode. Their recombinant nature is stably maintained because of a covalent junction between one of the viral genes and the full-size foreign gene inserted, a situation achieved here for the first time, via constructing an influenza ambisense RNA segment. The whole procedure is independent of any (selectable) phenotype, and can be applied to either of the eight influenza vRNA segments. After establishing a first ambisense vRNA segment carrying a single foreign gene it can also be repeated all over for inserting a second foreign gene within another ambisense RNA segment of the same constitution in principle.

Stable recombinant viruses of the type described can be used for cheap propagation in fertilized eggs, either for production of those recombinant viruses themselves or for production of foreign proteins or glycoproteins encoded by them, and hence find application in (glyco)protein production or in providing vector systems for somatic gene therapy or in being used as vaccination agents.

Thus, the present invention provides

(1) a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus, wherein

(a) at least one of the regular viral RNA segments has been exchanged for a vRNA encoding a foreign gene; and/or

(b) at least one of the regular viral RNA segments is an ambisense RNA molecule, containing one of the standard viral genes in sense orientation and a foreign, recombinant gene in anti-sense orientation, ~~or vice versa, in overall convergent arrangement;~~

(2) a method for the production of recombinant influenza viruses as defined in (1) above comprising

(a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense, sense or ambisense design,

(b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in at least one of its viral RNA segments, and

(c) thereafter selective vRNA inactivation through ribozyme cleavage;

(3) a method of constructing influenza carrier strains carrying one or more ribozyme target sites (of type one) in vRNA flanking positions comprising

(a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, carrying two different 3' promoter sequences in tandem, which are separated by a second type of ribozyme target sequence, and which carry the said internal ribozyme target sites of type one;

(b) followed by infection of an influenza wildtype strain;

(c) thereafter amplification through simple steps of viral propagation; and

(d) finally isolation through removal of their external 3' promoter sequence by ribozyme cleavage through infection of cells expressing ribozyme type 2, followed by plaque purification;

said method being suitable for the construction of an influenza carrier strain required for step

(b) of (2) above;

(4) a ribozyme-sensitive influenza carrier (helper) strain obtainable by the method of (3) above;

(5) a pharmaceutical composition comprising a recombinant virus as defined in (1) above;

(6) use of a recombinant virus as defined in (1) above for preparing a medicament for vaccination purposes;

- (7) use of a recombinant virus as defined in (1) above as vector systems in somatic gene therapy, for transfer and expression of foreign genes into cells (abortively) infected by such viruses, either in *ex vivo* or *in vivo* application schemes; and
- (8) a method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus as defined in (1) above as expression vector.

Moreover, the invention provides a method for preventing/treating influenza utilizing a recombinant virus as defined in (1) above, ~~i.e., a vaccination method utilizing said~~ recombinant virus.

According to the present invention "influenza virus" embraces influenza A virus, influenza B virus and influenza C virus. A preferred recombinant virus of the invention is where one or both of the standard glycoproteins hemagglutinin and neuraminidase have been exchanged into foreign glycoprotein(s) or into fusion glycoproteins consisting of an anchor segment derived from hemagglutinin and an ectodomain obtained from the foreign source, viral or cellular, or in which such recombinant glycoprotein has been inserted as a third molecular species in addition to the remaining standard components.

Further preferred recombinant viruses of the invention are where the terminal viral RNA sequences, which are active as the promoter signal, have been varied by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence.

Further preferred embodiments of the invention are set forth herein below.

A. Construction of influenza helper virus strains carrying ribozyme target sequences in flanking positions within either of the vRNA segments.

A.1. Influenza RNA polymerase will initiate transcription and replication from promoter structures located at internal positions in an RNA molecule, not only from the natural position at both ends of a vRNA molecule:



This is true in particular for promoter-up variants in RNA-internal location due to their enhanced binding affinity for viral polymerase. Not only 3' end extensions are tolerated in RNA-internal promoter recognition (Fig. 1), but also 5' extensions as well as extensions at both ends of the RNA template molecule, containing noncomplementary as well as complementary sequence, i.e. potentially present as a double-stranded extension. Finally, also an extension by way of duplication of the promoter sequence (active or inactivated) leads to mRNA transcription and CAT expression, initiated from the active pair of 5' and 3' promoter halves, irrespective if in external or in internal or even in an oblique localization (active 5' promoter sequence in external, active 3' promoter sequence in internal position, or vice versa). RACE-determination of the resulting 5' and 3' ends of viral mRNA and cRNA, i.e. the products of transcription and replication reactions for several of the extended template vRNA constructs proves an exact recognition and sequence-specific initiation at a position equivalent to regular 3' position 1: all of the various template extensions are lost in every product RNA molecule, most likely after only one round of replication.

A.2. Bicistronic (tandem) vRNA molecules carrying an additional 3' specific promoter sequence in a central position between its two genes can be used for an indirect selection method for recombinant influenza viruses:

The method described is applicable for any foreign gene (e.g. CAT) without a selection potential of its own, if inserted into the distal mRNA position (proximal vRNA position, in anti-sense orientation) behind a carrier gene (e.g. GFP) in the proximal mRNA position, able to serve as a transient selection marker. The carrier gene which is used for selection will get lost spontaneously during further propagation. These constructs are equivalent to a 3' extension of the template vRNA by a full-size gene of 750 nucleotides up to a second 3' promoter sequence, in terminal location. While in the set of experiments shown in Fig. 2 the external 3' promoter sequence is maintained throughout as the same promoter-up variant (1104), the internal 3' promoter sequence has been varied to include a full-level promoter-up variant (pHL2270, containing promoter mutant 1104), a medium-level promoter variant (pHL2350, promoter mutant 1948), a wildtype promoter construct (pHL2629), and a construct carrying an unrelated central sequence in an otherwise identical design (pHL2300).

Due to the presence of two 3' promoter sequences in conjunction with a single 5' promoter sequence an alternating interaction between them will constitute one or the other active

promoter structure (see Fig. 3). While the external location with an adjacent RNA 3' end may have a structural advantage, this appears to be compensated by the shorter distance in an interaction between the 5' sequence and the central 3' sequence in constituting the internal promoter, such that the competition between the two primarily reflects the various internal 3' promoter allele sequences used, compare Fig. 2B and activity ratios indicated above and below the lanes. Translation of the mRNA-distal gene (CAT) is only observed following an internal initiation at the bicistronic vRNA template, resulting in a spontaneous deletion of the

~~mRNA-proximal (vRNA-distal) gene, GFP, compare right half of Fig. 3 In complementary~~

analyses GFP fluorescence is observed initially for all of the bicistronic constructs, but gets lost on a faster rate from pHL2270 transfected and influenza infected cells (not shown), and will stay unchanged in pHL2300-treated cells. The indirect selection system based on bicistronic (tandem) molecules as designed here and demonstrated for reporter genes GFP and CAT can be used for any other gene without distinct phenotype upon insertion behind an unrelated carrier gene with properties useful in selection. – In employing that technique an initial phase of (repeated) positive selection for infected cells expressing that proximal trait (e.g. by FACS or by magneto-beads) will be followed by a second phase with negative selection, i.e. against that fraction of infected cells still exposing the same property.

A.3. Isolation of an influenza strain designed to carry 2x2 flanking ribozyme target sequences at the 5' and 3' end of vRNA segment 4 coding for hemagglutinin:

The above scheme for an indirect selection of any foreign recombinant gene behind a proximal carrier gene is further modified by deleting the carrier gene altogether. Instead, both 3' terminal promoter sequences (mutant and wildtype) follow each other at a short distance, separated only by a specific, repetitive ribozyme target sequence, - different from other target sequences to be described further below. The cDNA insert following after the second 3' promoter sequence consists mainly of a regular hemagglutinin (H7) coding sequence, however both the 5' and 3' vRNA terminal regions of the insert carry that other ribozyme target sequence (different from the first target sequence mentioned above) inserted in either location in tandem duplication (pHL2969, see Fig. 14).

Due to a superior replication supported by the promoter-up variant located in 3' external position the recombinant HA segment attached to that promoter sequence will become enriched during the first steps of viral propagation, while the originally dominating HA

segment of the helper virus which is under control of a wildtype promoter sequence is consecutively reduced and finally is no longer detectable among viral progeny. This result is documented by RT-PCR analysis of consecutive viral populations as obtained in that stepwise propagation procedure, see Fig. 4.

In the next step the viral lysate is twice passaged via infection of cell culture cells (293T) that before have been DNA-transfected by plasmid pAM403 (Fig. 15). This construct has been specifically designed to express a hammerhead ribozyme with flanking sequences

---

~~complementary to the repetitive GUC-containing target sequence, as present twice in between~~  
the external and internal 3' promoter signals in the recombinant HA vRNA segment, see Fig. 5. In this way the extra promoter sequence is cut off from the finally resulting recombinant HA vRNA. Its promoter-up activity was useful in achieving an initial increase in the concentration of recombinant HA vRNA over wildtype HA vRNA, and in finally excluding the latter from further propagation. However, for the same reason that high activity of the promoter variant will cause instability in the resulting viral progeny, and an effective 'substitution' at this time through ribozyme cleavage by the internally located promoter signal, wild-type or slightly improved, will restore stability to the progeny viruses, with all of their eight vRNA segments now brought back in balance to each other. Due to the ribozyme cleavage site at 26 nucleotides 3' of the wild-type promoter sequence (see Fig. 5), in the initial stage that promoter signal is situated in a vRNA-internal location, extended by a 3' terminal sequence of 26 nucleotides. According to the data presented in Fig. 1 this should cause a transient slight reduction in activity, resulting however in one step in regular viral mRNAs and cRNAs, with any initially remaining extra sequence being lost from the finally resulting recombinant HA vRNA.

Progeny viruses still carrying an external promoter-up sequence (before ribozyme treatment or due to incomplete reaction) will not cause any plaque, due to over-replication of one vRNA segment relative to all others which results in a high load of defective particles. However, progeny viruses which have lost that external promoter element due to ribozyme cleavage will yield regular plaques due to a balanced mode of replication for all eight wild-type or recombinant vRNA segments. Hence plaque purification is used for isolating a pure influenza viral strain carrying 2x2 ribozyme targets in its recombinant HA vRNA segment, with its termini reduced to the wild-type promoter sequence. The nature of the viral strain isolated has been confirmed in this regard by RT-PCR analysis, see Fig. 4.

The above isolation procedure resulting in influenza viral strains carrying 2x2 flanking ribozyme target sequences has been performed twice for the HA coding segment (segment 4) to obtain two different isolates with regard to the orientation of the ribozyme target sequences. In one of the isolates (vHM41, see SEQ. ID NO: 3) the tandem target sites have been inserted into the HA vRNA non-translated sequence both in 5' and 3' location, while according to the second design that 5' tandem target sequence has been included in an inverted orientation, such that it is now present in the cRNA 3' sequence instead (vHM42).

---

~~In another experiment the same procedure was used to isolate an influenza strain carrying 2x2~~  
tandem target sites within the 5' and 3' flanking positions of segment 8 vRNA, i.e. coding for genes NS1 and NS2 (vHM81; see SEQ.ID NO:4). And in principle the same could be done for any other influenza segment, in particular since only the reading frame cDNA sequence has been exchanged from HA to NS, with all of the flanking elements directly responsible for that procedure remaining in place, unchanged.

A.4. Ribozyme cleavage and vRNA segment exchange using ribozyme-sensitive influenza strains as helper viruses:

In an initial model experiment a range of ribozyme type and target site locations was probed in designing a series of CAT reporter gene vRNA constructs (analysed in the presence of a surplus of wildtype helper virus) in 293T cells. While all of the ribozyme constructs adhered to the hammerhead model, with 10 to 12 nucleotides of complementary sequence flanking on either side of the GUC target site, these ribozyme constructs varied from monomer to dimer to trimer repetitions. Ribozyme containing mRNAs were synthesized *in vivo* via the basic vector plasmid pSV2-*neo*, i.e. using the efficient p<sub>SV</sub> RNA polymerase II promoter element for expression, and the SV40 origin signal for plasmid amplification, in a cell line (293T or cos-1) with an incorporated SV40 T antigen gene. In addition the pSV2-*neo* mRNA includes the small, 63 nucleotide intron sequence of the SV40 early mRNA which is supposed to be spliced very slowly, thereby extending the pre-mRNA half-live in the nucleus. Each of the pSV2-*neo*-ribozyme plasmid constructs was transfected into 293T cells. Thereafter, recombinant viruses containing dimer target sites either near one end of the molecule only, or near both ends have been used for infection of the transfected cells. Relative activities of ribozyme constructs versus vRNA target sites have been measured via inactivation of CAT acetyl transfer rates in the cell lysates obtained at 8 h post infection (Fig. 6). The highest

activities were observed for dimer ribozymes acting on vRNA molecules carrying 2x2 target sequences on both ends of the molecule, either in vRNA 5' and 3' location, or in vRNA 3' and cRNA 3' location, i.e. with an inversion of the target site sequence at the vRNA 5' end.

Consequently, the two constructs carrying tandem ribozyme double targets within both of their non-translated vRNA flanking sequences have been used in the design of ribozyme-sensitive influenza virus strains as described above, with both variants isolated for segment 4 (HA), and one of them for segment 8 (NS). In complementary correspondence the

---

~~hammerhead ribozyme plasmid used has also been constructed as a double headed structure~~  
with flanking sequences as shown in Fig. 7 (pAM424; for its detailed structure, see Fig. 16).

The three target site-containing influenza strains isolated as described above have been analysed for their sensitivity against ribozyme cleavage by infection of 293T cells, which had been DNA-transfected 18 h earlier by ribozyme-producing plasmid pAM424, at DNA-transfection rates between 60 and 70 % (as observed in parallel transfections using GFP-expressing plasmid pAM505). Inactivation rates in these one-step control experiments were between 90% and 99% for all three ribozyme-sensitive strains, in their extent mainly depending upon the actual DNA-transfection rates achieved in individual experiments.

In the next step both isolates of HA-coding ribozyme-sensitive viruses, vHM41 and vHM42, have been used in marker-rescue experiments. Here, 293T cells have been first DNA transfected by HA-variant cDNA construct pHL2507 (see Fig. 17), followed after 18 h by vHM41 or vHM42 virus infections at moi 1. The resulting viral supernatant containing e.g. a mixture of ribozyme-sensitive vHM41 carrier virus and pHL2507/vHM41 recombinant virus is propagated in an intermediate step on MDCK cells, which also results in an increase in the fraction of recombinant viruses. Thereafter the resulting virus-containing supernatant is passaged through 293T (or cos-1) cells, which in advance have been transiently transfected by ribozyme-producing pAM424. As may be concluded from the above experiment (Fig. 6) and shown in Fig. 8 vHM41- or vHM42-derived ribozyme-sensitive HA vRNA segments are expected to be inactivated by pAM424-produced ribozyme down to a remaining level of about 1% to 10% (mainly present within cells that are infected, but not DNA-transfected).

Instead, the substitute HA vRNA which originated from pHL2507 plasmid DNA (ribozyme-resistant) becomes exclusively incorporated into progeny virions. For further purification and

viral propagation these have been passaged a second time through 293T cells, again in advance DNA-transfected by pAM424, and after another amplification step on MDCK cells the resulting viral preparations have been used for RT-PCR analysis. The resulting viral populations in these marker rescue experiments only contain HA-vRNA molecules derived from pHL2507, which in their PCR analyses are of intermediate size relative to wildtype HA-vRNA, and vHM41- or vHM42-derived HA-vRNAs, respectively.

Consequently, a set of ribozyme-sensitive influenza strains with targets inserted individually into every vRNA molecule may be used for such one-step marker rescue experiments in general, i.e. for vRNA segment exchange reactions performed in a directional way for any of the eight influenza vRNA segments, without requirement for a selectable change in phenotype (genetic marker).

#### B. Expression of two gene products from ambisense bicistronic influenza vRNA

B.1. The influenza cRNA promoter is active in antisense viral mRNA transcription according to the cap-snatching mode of initiation:

While the vRNA template of influenza virus is known to be active in viral mRNA as well as cRNA synthesis, the cRNA template has been described so far only to produce vRNA molecules, as a second step in viral replication. The potential activity of the cRNA promoter in initiating also viral mRNA transcription has not been analysed or even suspected so far, since no antisense (vRNA) reading frame can be detected in any of the viral RNA segments. Also, no  $U_3/U_6$  template sequence element is present in any of the viral cRNAs in an adjacent position to its 5' promoter structure as is the case for all viral vRNAs. This element is known to serve as a template sequence for mRNA terminal poly-adenylation, in repetitive interaction. However, when both elements are provided through reconstruction of an artificial influenza cRNA segment: a reading frame in opposite orientation (CAT), and a  $U_6$  template element in 5' adjacent location, CAT expression can indeed be observed, see pHL2583 (see Fig. 18) in Fig. 9. Similar to the vRNA promoter the cRNA promoter activity is improved by (the same) promoter-up mutations, which essentially consist of basepair exchanges according to the 'corkscrew' model. This model apparently also holds for the cRNA promoter structure as analysed in a stepwise manner in Fig. 9. While the cRNA promoter has to be superior over the vRNA promoter in its initiation of replication, since the vRNA/cRNA product ratio was

determined to be around 10:1 (Yamanaka et al., *In vivo* analysis of the promoter structure of the influenza virus RNA genome using a transfection system with an engineered RNA. Proc. Natl. Acad. Sci. USA 88, 5369-5373 (1991)), the cRNA promoter is observed to be inferior to the vRNA promoter in its initiation rate of transcription (compare pHL2583 with pHL1844 in Fig. 9), at least for all promoter variants tested so far.

A RACE analysis for determination of the 5' ends in pHL2583 cRNA promoter transcribed mRNAs proved this initiation to occur according to the cap-snatching mode, in complete equivalence to standard vRNA promoter-controlled transcription initiation.

---

## B.2. Development of ambisense influenza constructs for consecutive expression of two genes (GFP and CAT) from a single viral RNA:

For a bidirectional transcription and translation of influenza RNA segments the two reporter genes GFP and CAT have been arranged in opposite orientation to each other, and the flanking 5' and 3' promoter sequences (adhering to promoter-up variant 1104) had to be reconstructed to include a U<sub>6</sub> poly-adenylation element in either orientation in a 5' promoter adjacent position. This requirement necessarily resulted in a promoter-adjacent 5'-U<sub>6</sub>/3'-A<sub>6</sub> complementary structure, both in the vRNA and cRNA terminal sequence (see Fig. 10), which had to be tested for its promoter activity, in either orientation. Therefore, the convergent pair of reporter genes GFP and CAT has been inserted in both orientations, such that CAT transcription is initiated by the vRNA promoter in one construct (pHL2960), and by the cRNA promoter in the other (pHL2989, Fig. 19), and vice versa for GFP expression from both ambisense constructs. In addition, also the CAT gene only has been inserted in either orientation between the 5' and 3' elements of that ambisense promoter, with CAT transcribed by the vRNA promoter in one case (pHL2959), and by the cRNA promoter in the other (pHL2957). The whole set of constructs allows for a direct comparison with corresponding reference constructs carrying a regular vRNA promoter (pHL1844) or cRNA promoter structure (pHL2583), i.e. carrying only the 5'-adjacent U<sub>6</sub> sequence element and no 3'-A<sub>6</sub> counterpart. The two groups of constructs also differ in insert size, since a single inserted gene roughly accounts for 750 nucleotides, the convergent set of two genes for 1500 nucleotides, with the distal half of both mRNAs in this case remaining untranslated, a situation unusual for influenza viral mRNAs.



As is demonstrated in Fig. 10B for CAT expression of the various ambisense constructs all of them are able to initiate transcription in both orientations, even if at different levels with regard to their vRNA and cRNA promoter-dependent expressions, and also with regard to the insert lengths and convergent arrangements of the GFP/CAT versus CAT-only constructs. Analysis of the GFP expression rates (not shown) yields complementary results, i.e. again vRNA promoter-controlled GFP expression is superior over cRNA promoter expression of GFP. Therefore, individual ambisense clones either show an asymmetric high expression of ~~GFP and low expression of CAT (pHL2960) or vice versa (pHL2989), depending on their~~ orientation of reading frames with regard to the external vRNA and cRNA promoter. Fig. 10C also demonstrates successful propagation of recombinant viruses containing ambisense RNA molecules, which proves survival through amplification, packaging into virions, and expression of both mRNAs in infected MDCK cells (including besides CAT also GFP expression).

B.3. Construction of a superior promoter-up mutation, pHL1920, to be used for improved rates of cRNA promoter expression in ambisense constructs:

An extended analysis of promoter variants, in particular of complementary double exchanges according to the 'corkscrew' model yielded among others variant pHL1920 (Fig. 20) with CAT activity rates considerably above (125-130% of) the rates observed for standard promoter-up variant '1104' (as present in pHL1844). The '1920' promoter-up variant consists of altogether 5 nucleotide substitutions relative to the wildtype promoter sequence, both in the 5' promoter element (2), and the 3' promoter element (3). The structure of this variant and the whole set of complementary double exchanges is presented in Fig. 11, together with the respective CAT activity measurements, in vRNA promoter constructs. vRNA promoter-up variants also show similarly improved expression in (ambisense) cRNA constructs, even if at generally lower levels than in vRNA constructs. cRNA promoter-up expression is observed at levels similar or somewhat (2x-5x) above the *wild-type* vRNA promoter rate, while vRNA promoter-up constructs show CAT expression rates increased up to 20 or 25 times the wild-type vRNA promoter level. In either case expression rates also depend on the size of the insert, with promoter activity rates decreasing with increasing lengths of the influenza RNA molecules to be transcribed.



B.4. Influenza recombinant viruses containing a foreign gene (CAT) in covalent ambisense linkage with one of the viral genes (HA, NS1/NS2):

The principle solution in designing stable recombinant viruses based on the new properties observed for influenza transcription and replication signals consists in constructing viruses which contain a foreign gene in covalent linkage with one of the (indispensable) viral genes, in ambisense bicistronic organization. Preferably the viral gene is connected to the cRNA promoter, while vRNA promoter expression is used for expression of the foreign gene at rates

---

considerably above the viral mRNA synthesis. The promoter-up variant chosen for constructing the ambisense RNA segment intends to bring its cRNA promoter expression (approximately) into balance with all other viral gene expression levels, which are controlled by wild-type vRNA promoters located at the termini of the seven ordinary influenza segments; the respective choice has to take into consideration the overall length of the ambisense segment.

Isolation of the ambisense recombinant virus employs an RNA polymerase I-transcribed ambisense cDNA construct, which will give rise *in vivo* to ambisense cRNA-type molecules, see Fig. 12. The plasmid DNA transfection mixture used in this step with 293T cells in addition may or may not contain four 'booster' plasmids which under p<sub>CMV</sub>-control produce the four early influenza proteins from non-viral mRNAs: NP, plus PB1, PB2, and PA, i.e. the three subunits of viral polymerase (Pleschka et al., A plasmid-based reverse genetics system for influenza A virus J. Virol. 70, 4188-4192 (1996)), which will increase in a pre-amplification step the copy number of that ambisense viral cRNA segment. At 18 h post transfection the 293T cells are infected by a ribozyme-sensitive influenza strain, e.g. vHM41, which will supply (again) early and also late viral RNAs. The resulting supernatant which contains a mixture of vHM41 carrier virus and vHM41-derived ambisense recombinant virus (nine vRNA segments) is then passaged directly or via an intermediate step of amplification on MDCK cells onto 293T cells that have in advance been DNA-transfected by ribozyme-producing pAM424. Here, the ribozyme-sensitive vRNA segment of vHM41 will be cleaved at its 2x2 target sites by pAM424 specific ribozymes. In recombinant viruses the vRNA gene lost in this way is re-supplemented through its presence within the ambisense segment. The virus-containing supernatant is passaged for amplification and further purification through ribozyme treatment a second time on 293T cells which again have been pretreated by pAM424 DNA transfection.

Absence of ribozyme-sensitive vRNA, and presence only of ambisense RNA in RT-PCR analysis at this stage allows for further amplification on MDCK cells and a final virus stock preparation on embryonated chicken eggs. CAT assays can be used to analyse for the presence and monitor the activity of this model foreign gene through the various steps of isolation and propagation as well as document technical improvements that might be worked out for one or more of the processive stages.

---

C. Examples for application of helper-free, stable recombinant influenza viruses

C.1. Incorporation of reporter gene GFP in NS/GFP or HA/GFP ambisense segments:

Recombinant viruses of this type will allow to follow-up on influenza infection instantly and continuously in individual infected cells, which may also be counted or documented by fluorescence photography or FACS sorting. With improved temperature resistance (Siemering et al., Mutations that suppress the thermosensitivity of green fluorescent protein. *Curr. Biol.* 6, 1653-1663 (1993)) GFP expression becomes visible at 2 h after infection and shows bright fluorescence after 4 h p.i., it will be possible to follow-up on the spread of viral infection by GFP fluorescence. Stable fluorescence in abortively infected cells e. g. observed in *ex vivo* treatment of dendritic cells similarly supports a follow-up on their reincorporation into animals; other genes may be incorporated by ambisense vRNA into dendritic cells in the same way.

C.2. Construction of glycoprotein CSFV-E2 carrying influenza, helper-free:

The glycoprotein E2 of CSF virus has been incorporated previously into influenza both as an HA-anchor fusion protein within the viral envelope, and as an additional, unstable ninth vRNA segment into its genome (Zhou et al.; Membrane-anchored incorporation of a foreign protein in recombinant influenza virions. *Virology*, 246, 83-94 (1998)). Stabilization is now achieved through an ambisense connection with either of the regular viral RNA segments (NS or HA) which also allowed to reach a level of 100% recombinant viruses instead of an hitherto only 20% (Fig. 13), since all carrier viruses are destroyed through ribozyme action, see Figs. 8 and 12. The helper virus containing preparation has already been used successfully as a vaccine against CSFV infection (antibody titers of 1:40000); the increase achieved in recombinant viruses allows a further improvement in that regard. Also (cost-

effective) propagation in fertilized chicken eggs has become possible due to its stable incorporation of the foreign gene as a covalent ambisense construct.

C.3. Construction of hepatitis C glycoprotein-carrying recombinant influenza viruses as a candidate vaccine:

Hepatitis C virus is a close relative of CSF virus (hog cholera virus), and in particular its set of two glycoproteins, small-size E1 and larger-size E2, is closely related in structural detail and presumably also in function to the corresponding CSFV proteins. An incorporation of HCV-E2/HA fusion proteins into influenza viral envelopes has been achieved in analogy to the CSFV-E2/HA incorporation. In addition, incorporation of an anchor-fusion glycoprotein HCV-E1/HA or both together (in NS *and* HA ambisense junctions) allows further variations in constructing an influenza-based vaccine for hepatitis C. In analogy to CSFV-E2, neutralizing antibodies are expected to be directed against particular epitopes of HCV-E2, presented in essentially native conformation at the influenza viral envelope.

C.4. Stable incorporation of selected influenza T-cell epitopes in ambisense constructions:

Influenza infection is known to result in both, antibody production against that specific viral strain or indeed its epitopes that are located mainly at the surface of HA, and in an increase of specifically primed cytotoxic T-lymphocytes, stimulated by T-cell epitopes primarily located within the essentially invariable core structure of the NP protein. While the humoral response will result in life-long immunity against *that particular* strain of influenza or its epitope structures, the T-cell response will be lost or severely reduced some time afterwards, such that its specificity against influenza in general will fall below protective levels. One way in trying to increase that level of cellular immunity is to enhance the response or recruitment of influenza-specific CTL cells by increasing the level of T-cell epitopes in the infected cell and hence its presentation on the surface by MHC-I receptors. This is achieved by combining in an ambisense construct the HA gene and a series of repeated T-cell epitope sequences as present in the influenza NP gene, in a model design specific for the BALB/C mouse MHC-I allele. Here, the promoter-up expression rate is realized (in vRNA promoter-controlled initiation) for expression of the repetitive epitope polypeptide chain. Alternatively or in addition a controlled secretion of an interleukin can be achieved from recombinant influenza-infected cells, upon ambisense incorporation of the respective gene preferably into the NS

segment. The interleukin to be chosen for this purpose (IL-12 or other) is selected to enhance the longevity of influenza-specific CTL cells or its conversion into corresponding memory cells. In this way an ambisense vaccine strain against influenza itself is achieved with expected protective capacity against influenza in general.

C.5. Exchange of influenza glycoproteins against foreign viral glycoproteins (VSV-G):

The 'marker rescue' experiment described above (section A.4.), i.e. an exchange of one HA gene (ribozyme-sensitive) for another (ribozyme-resistant) can also be used for an exchange of HA for an entirely different glycoprotein, such as the vesicular stomatitis virus G protein, as long as it is attached to the HA anchor segment. Effective incorporation depends always on that C-terminal tail sequence and its interaction with underlying matrix protein M1, and therefore, all constructions consist of fusion proteins in direct analogy to CSFV-E2. VSV-G with or without a foreign anchor sequence has been shown in several other viruses to be able to substitute for the original glycoprotein and to result in infectious viruses with VSV-G specific host-ranges (e.g. in retroviruses, rabies virus, measles virus). G protein in VSV itself as well as on the surface of foreign viruses is the only glycoprotein required for all of the consecutive steps in infection. Insertion of VSV-G instead of HA in recombinant influenza viruses leaves the second glycoprotein, neuraminidase, without any function, which then will get lost spontaneously from the recombinant viruses. This will further increase the capacity for an addition of foreign genes, beyond the gain resulting from an exchange of the larger HA for the smaller VSV-G, which might be used for an addition of ambisense constructs.

The invention is further illustrated in the accompanying figures.

**Description of the figures**

**Fig.1. 3' nucleotide extensions of influenza vRNA template molecules:**

(A) Murine B82 cells have been transfected by plasmid cDNA constructs designed to be transcribed into influenza vRNA molecules by RNA polymerase I *in vivo*, followed after 20 h by standard FPV<sub>Bratislava</sub> helper virus infection, at an moi of 1 to 3. In addition to reference plasmid pHL2024 (no extension), related cDNA constructs carrying extensions of 1 to 50 bp, and hence extended by 1 to 50 nucleotides at the resulting vRNA 3' ends were used in parallel

transfections; template extensions are marked at the top of the figure. Cell lysates prepared at 8 h post helper virus infections were used for CAT reactions using in one round 50  $\mu$ l of cell lysate each, and in further analyses 5  $\mu$ l and 0.5  $\mu$ l of lysate (not shown). Relative yields were determined in comparison to reference plasmid pHL2024, as indicated below the figure, with calculations restricted to those CAT assays showing less than 40% of substrate consumption, in three or more independent experiments.

(B) Viral passage of B82 supernatants containing recombinant influenza virus onto MDCK cells, at an moi of 2 to 4, in average. Again at 8 h post infection cell lysates have been prepared and used for CAT assays. Relative yields as indicated below the figure have been determined in comparison to pHL2024 the same way as in (A), using 0.5  $\mu$ l of cell lysate in each case. (The 50  $\mu$ l CAT assays as shown here and also in the following figures intend to give an immediate impression of relative activities at always the same level, while the actual measurement data as indicated below the lanes are obtained at various appropriate enzyme concentrations relative to reference pHL2024.)

Fig. 2. Propagation of recombinant influenza viruses with tandem bicistronic vRNA:

(A) General design of expression plasmids for transient bicistronic vRNAs coding for GFP in the mRNA-proximal, and for CAT in the mRNA-distal position. Among the functional elements indicated are the human RNA polymerase I promoter ( $p_h$ ) and murine rDNA terminator ( $t_r$ ) sequences, both hatched, and the 5' and 3' vRNA promoter cDNA sequences, open and closed boxes, respectively. For the vRNA-internal 3' promoter signal three variant sequences have been inserted as indicated below (positions 1 to 15 refer to 3'-terminal nucleotides in the resulting monocistronic vRNAs).

(B) CAT assays as determined relative to pHL1844 (monocistronic CAT construct) after DNA transfection of 293T cells plus helper virus infection followed by one round of progeny viral propagation on MDCK cells are indicated *below* the lanes. Relative activities of the internal promoter sequences as indicated *above* the figure refer to measurements in a monocistronic *external* location of the same promoter variants (Flick *et al.*, Promoter elements in the influenza vRNA terminal structure, RNA 2, 1046-1057 (1996)). Control clone pHL2300 contains an unrelated, non-functional sequence in the central location in an otherwise identical plasmid construct.

Fig. 3. Tandem bicistronic vRNA sup-orting an alternative mode of tran-scription and replication initiation:

An additional internal 3' promoter sequence has been inserted in between both cistrons, in a vRNA-central position. Left half: bicistronic replication and transcription leading to (proximal) GFP expression. Right half: internal initiation resulting in monocistronic replication and transcription leading to (distal) CAT expression, and causing deletion of the GFP sequence from progeny molecules.

Fig. 4. Outgrowth of promoter-up recombinant vRNA versus wildtype vRNA segments in stepwise propagation of influenza virus:

RNA polymerase I transcription of transfected pHL2969 DNA results in influenza vRNA carrying (an external) promoter-up mutant '1104', and containing 2x2 ribozyme targets in flanking positions relative to its HA coding sequence. Another HA vRNA segment in wild-type configuration and originating from infecting FPV helper virus is also present in the recombinant virus preparation, initially (lane 1; 293T lysate) in surplus amounts, but reduced and finally lost entirely in consecutive steps of propagation (lanes 2 to 4; MDCK cell lysates), and in isolated strains after pAM403 ribozyme treatment for removal of the external '1104' promoter sequence (lane 5). Determination throughout by RT-PCR analyses using a pair of primers extending across the 5' inserted target site sequence, with 435 bp representing the recombinant HA segment, and 306 bp the wild-type sequence without an inserted target site sequence.

Fig.5. pAM403 ribozyme cleavage of pHL2969 derived vRNA molecules at specific target sites inserted between an external and an internal 3' promoter sequence:

The external promoter-up ('1104') signal is used for vRNA amplification within recombinant viruses and reduction of helper virus HA vRNA (Fig. 4), while the 'switch' to an internal wild-type signal guarantees stable replication of recombinant viruses. pAM403 hammerhead ribozyme RNAs are indicated in complementary binding to their target site sequences (12 and 10 nucleotides flanking the GUC cleavage point) by straight lines flanking a central secondary structure symbol. vRNA-internal 2x2 ribozyme targets are marked by xx (see Fig. 7).

Fig. 6. Comparative cleavage analysis of model CAT vRNAs with tandem target sites in various flanking positions, by target-specific ribozymes:

293T tissue culture cells have been transiently DNA-transfected either by a single-headed hammerhead ribozyme (s), or a double-headed (d), or triple-headed (t) ribozyme cDNA construct, all specifically designed to hybridize to a tandem dimer target site sequence inserted in flanking positions into the CAT vRNA. All ribozyme RNAs have been expressed ~~from the same pSV2-neo plasmid vector, including a pSV2-neo control construct without an~~ inserted ribozyme cDNA sequence (c). At 20 h after DNA transfection (which reached 65% yield as measured by  $P_{CMV}$ -GFP transfection in parallel of the same cell culture) the 293T cells were infected by CAT recombinant viruses carrying tandem double target sequences either only in vRNA-3' position, or in both vRNA-3' and 5' positions, or in both vRNA-3' and cRNA-3' positions. Most effective among the s, d, or t-ribozymes were double-headed constructs, acting on 2x2 targets inserted in either of the two localizations described (lanes 6 and 10).

Fig. 7. Alignment of pAM424 double-headed ribozyme with one of their repetitive target sequences located within the 5' and 3' vRNA flanking regions:

The superior activity of ribozymes oriented against targets located in the 3' end of vRNA molecules over those present in the 5' end instead (not shown) is in agreement with the model for influenza vRNA transcription and replication (Lamb and Krug, Orthomyxoviridae: The viruses and their replication. In 'Virology' (B.N.Fields, D.M.Knipe, P.M.Howley, R.M.Chanock, J.L.Melnick, T.P.Monath, B.Roizman, and S.E.Straus, Eds.), 3<sup>rd</sup> ed., Vol. 1, pp. 1353-1395. Lippincott-Raven, Philadelphia (1996)), according to which influenza polymerase stays attached to the 5' end of the vRNA molecule throughout the entire or even several rounds of transcription, whereas the very 3' end repeatedly, in every initiation reaction serves as the template sequence, and consequently is no longer covered by polymerase. Superiority of a double-headed over a single-headed ribozyme has been determined earlier in this laboratory (A.Menke, Anti-Influenza Ribozyme: vRNA-Spaltung und intrazelluläre Aktivität. Dissertation Universität Giessen (1997)), but the substantial increase of vRNA inactivation rates upon incorporation of tandem target sites at both ends of the vRNA

molecule instead of only one has been observed here for the first time, within that overall design.

Fig. 8: pAM424 ribozyme cleavage of resistant FPV wild-type HA vRNA and ribozyme-sensitive PHL2969-derived HA-vRNA in 293T cells infected by vHM41 after isolation from PHL2969-recombinant viral preparations. Lane 1: FPV infection of 293T cells, untreated; lane 2: FPV infection of 293T cells DNA-transfected by pAM424; lane 3: vHM41 infection of 293T cells, untreated; lane 4: vHM41 infection of 293T cells DNA-transfected by pAM424. RT-PCR analyses of purified viral progeny as in Fig. 4.

Fig. 9 Functional analysis of the influenza cRNA promoter structure:

(A) Schematic cRNA promoter ('1104') secondary structure according to the 'corkscrew' model; nucleotides involved in single or double nucleotide exchange are marked by their position.

(B) CAT analyses of 293T cell lysates after DNA transfection and FPV helper virus infection of cRNA promoter variants, in comparison to standard vRNA promoter-up mutant '1104' (PHL1844). Nucleotide substitutions divergent from the basic '1104' structure as present in PHL2583 or PHL2721 (see above) are indicated above the lanes, positions  $\bar{3}$  or  $\bar{8}$  as marked by a bar refer to cRNA positions counted from the 3' end. Relative CAT activities are marked below the lanes.

Fig.10 Functional analysis of the vRNA and cRNA promoter in ambisense arrangement:

(A) Sequence organisation of the ambisense promoter cDNA construct carrying  $T_0/A_0$  elements adjacent to the terminal sequence, and secondary structure predictions for the resulting cRNA and vRNA promoter signal.

(B) CAT expression data obtained from the cell lysates of 293T cell after plasmid DNA transfection and FPV infection, and (C) from cell lysates of MDCK cells after one step of viral passage. Indicated above the lanes are promoter/gene conjunctions: v = vRNA promoter; c = cRNA promoter.



Fig.11 Basepair substitutions according to the vRNA 'corkscrew' structure:

(A) 'Corkscrew' conformation of the vRNA promoter drawn against a schematic indication of interacting tripartite viral polymerase. Paired positions exchanged in individual experiments are indicated by numbers, nucleotides  $\bar{3}$  or  $\bar{8}$  are counted from the 3' end. pHL2024 containing promoter-up mutation '1104' is used as the reference construct (=100%) in all of the CAT assays, while pHL2428 represents the wild-type promoter structure.

(B) CAT expression data obtained after one step of viral passage in MDCK undiluted, and 50 fold diluted.

Fig. 12: Flow-chart of the isolation procedure for an ambisense recombinant influenza virus.

Fig. 13 Immuno-electron microscopy of purified influenza FPV/CSFV-E2-HA virions:

Recombinant viruses exposing the foreign glycoprotein CSFV-E2 in their envelopes, which has been fused onto the HA anchor domain, are marked by anti-E2 monospecific antibody and by secondary gold-labelled (5nm) goat antibody. Recombinant viruses (16%) are present together with their FPV helper viruses.

Fig. 14: pHL2969; the exact sequence of the 4930 bps circular DNA is shown in SEQ. ID NO:1.

Fig. 15: pAM403; the exact sequence of the 5811 bps circular DNA is shown in SEQ. ID NO:2.

Fig. 16: pAM424; the exact sequence of the 5860 bps circular DNA is shown in SEQ. ID NO:5.

Fig. 17: pHL2507; the exact sequence of the 4610 bps circular DNA is shown in SEQ. ID NO:6.

Fig. 18: pHL2583; the exact sequence of the 3558 bps circular DNA is shown in SEQ. ID NO:7.

Fig. 19: pHL2989; the exact sequence of the 4343 bps circular DNA is shown in SEQ. ID NO:8.

Fig. 20: pHL1920; the exact sequence of the 3888 bps circular DNA is shown in SEQ. ID NO:9.

Hence, the present invention is based on two surprising findings, namely

1. influenza virus promoters are active when present internally in a gene;

2. the so-called cRNA, thought to be an intermediate in replication can be turned into a protein-encoding RNA by equipping it with a variant influenza virus promoter, described in the present invention.

These two observations were used to make ambisense constructs. This allows to package an additional, foreign gene into influenza virus particles. Such particles were made previously, by other methods, but proved to be unstable, and therefore useless. For use as a vaccine for example, a helper virus would have been needed as a stabilizer. Stabilization in the present invention is achieved by several means. These include the "balancing" of one of the two promoters in the ambisense biscistronic genetic construct with seven other vRNA wildtype promoters, while the additional promoter is used for high-rate expression of the foreign gene at various levels.

Thus, the present invention provides a system for expression of foreign proteins in higher eukaryotic systems. One system in particular is interesting, namely embryonated chicken eggs, as it allows cost-effective production in an automatable way (as used by most flu vaccine producers). The reason that this process is now possible, is that the foreign protein is part of a stable, engineered influenza virus particle. The virus can be designed also to rapidly monitor process improvements.

An excellent use is of course the use of the construct as a vaccine. The influenza virus particle is immunogenic and can now be equipped with foreign antigens, enabling for example the design and production of hepatitis C virus and HIV vaccines, but also of tumor vaccines. As the present invention shows, the foreign antigenic surface glycoprotein is "fused" to a C-

terminal segment of influenza HA, and the antigen then is presented at the surface of influenza virus particles. In addition, these vaccines can now be made in the way standard flu vaccines are made, i.e., in embryonated chicken eggs.

---

EPO - Munich  
33  
06. März 1999

&lt;110&gt; ARTEMIS PHARMACEUTICALS GmbH

<120> Stable Recombinant Influenza Viruses Free of Helper  
Viruses

&lt;130&gt; 990383ep

&lt;140&gt;

&lt;141&gt;

&lt;160&gt; 9

&lt;170&gt; PatentIn Ver. 2.1

&lt;210&gt; 1

&lt;211&gt; 4930

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 1

cgtacgaagc	ttctagaggg	attggctgag	acgaaaaaca	tatgctagag	ggattggctg	60
agacgaaaaa	catatgctag	agcggccgcc	accgcggtgg	agctccagct	tttgttccct	120
ttagtgaggg	ttaattgcgc	gcaggcctag	ctaggtaaaag	aaaaataccc	ttgtttctac	180
taataacccg	gcgGCCcaaa	atgccgactc	ggagcgaaaag	atatacctcc	cccggggccg	240
ggaggtcgcg	tcaccgacca	cgccgcgggc	ccaggcgacg	cgcgacacgg	acacctgtcc	300
ccaaaaacgc	caccatcgca	gccacacacg	gagcgcccg	ggccctctgg	tcaaccccag	360
gacacacgcg	ggagcagcgc	cgggcggggg	acgccctccc	ggccgcccgt	gccacacgca	420
gggggcccgc	ccgtgtctcc	agagcgggag	ccggaagcat	tttcggccgg	cccctcctac	480
gaccgggaca	cacgagggac	cgaaggccgg	ccaggcgca	cctctcgggc	cgcacgcgcg	540
ctcagggagc	gctctccgac	tccgcacggg	gactcgccag	aaaggatcgt	gacctgcatt	600
aatgaatcag	gggataacgc	aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	660
aaccgtaaaa	aggccgcgtt	gctggcggtt	ttccataggc	tccgcccccc	tgacgagcat	720
cacaaaaatc	gacgctcaag	tcagagggtg	cgaaccgccg	caggactata	aagataccag	780
gcgtttcccc	ctggaagctc	cctcgtgcgc	tctcctgttc	cgacctgtcc	gcttaccgga	840
tacctgtccg	cctttctccc	ttcgggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	900
tatctcagtt	cggtgtaggt	cgttcgctcc	aagctgggct	gtgtgcacga	acccccggtt	960
cagcccagcc	gctgcgcctt	atccggtaac	tatcgtcttg	agtccaaccc	ggtaagacac	1020
gacttatcgc	cactggcagc	agccactggg	aacaggatta	gcagagcgag	gtatgtaggc	1080
ggtgctacag	agttcttgaa	gtggtggcct	aactacggct	acactagaag	gacagtattt	1140
ggtatctgcg	ctctgctgaa	gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	1200
ggcaaaacaa	ccaccgctgg	tagcgggtgg	ttttttgttt	gcaagcagca	gattacgcgc	1260
agaaaaaaag	gatctcaaga	agatcctttg	atctttttcta	cggggtctga	cgctcagtgg	1320
aacgaaaact	cacgttaagg	gattttggtc	atgagattat	caaaaaggat	cttcacctag	1380
atccttttaa	attaaaaatg	aagttttaaa	tcaatctaaa	gtatatatga	gtaaacttgg	1440
tctgacagtt	accaatgctt	aatcagtgag	gcacctatct	cagcgatctg	tctatttcgt	1500
tcattccatag	ttgcctgact	ccccgtcgtg	tagataacta	cgatacggga	gggcttacca	1560
tctggcccca	gtgctgcaat	gataccgcga	gacccacgct	caccggctcc	agatttatca	1620
gcaataaacc	agccagccgg	aagggccgag	cgcagaagtg	gtcctgcaac	tttatccgcc	1680
tccatccagt	ctattaattg	ttgccgggaa	gctagagtaa	gtagttcgcc	agttaaatag	1740
ttgcgcaacg	ttgttgccat	tgctacaggc	atcgtgggtg	cacgctcgct	gtttggtagt	1800
gcttcattca	gctccgggtc	ccaacgatca	aggcgagtta	catgatcccc	catgttggtg	1860
aaaaaagcgg	ttagctcctt	cggtcctccg	atcgttgcta	gaagtaagtt	ggccgcagtg	1920
ttatcactca	tggttatggc	agcactgcat	aattctctta	ctgtcatgcc	atccgtaaga	1980

27

tgcttttctg	tgactgggtga	gtactcaacc	aagtcattct	gagaatagtg	tatgcggcga	2040
ccgagttgct	cttgcccggc	gtcaacacgg	gataataccg	cgccacatag	cagaacttta	2100
aaagtgtca	tcattggaaa	acgttcttcg	gggcgaaaac	tctcaaggat	cttaccgctg	2160
ttgagatcca	gttcgatgta	acccactcgt	gcaccaact	gatcttcagc	atcttttact	2220
ttcaccagcg	tttctgggtg	agcaaaaaca	ggaaggcaaa	atgccgcaaa	aaagggaata	2280
agggcgacac	ggaaatgttg	aatactcata	ctcttccttt	ttcaatatta	ttgaagcatt	2340
tatcaggggt	attgtctcat	gagcggatac	atatttgaat	gtatttagaa	aaataaaca	2400
aagagtttgt	agaaacgcaa	aaaggccatc	cgtcaggatg	gccttctgct	taatttgatg	2460
cctggcagtt	tatggcgggc	gtcctgcccg	ccaccctccg	ggcgttgct	tcgcaacgtt	2520
caaaccgct	cccggcggat	ttgtcctact	caggagagcg	ttcaccgaca	aacaacagat	2580
aaaacgaaag	gcccagtcct	tcgactgagc	ctttcgtttt	atttgatgcc	tggcagttcc	2640
ctactctcgc	atggggagac	cccacactac	catcggcgct	acggcgtttc	acttctgagt	2700
tcggcatggg	gtcaggtggg	accaccgcgc	tactgccgcg	aggcaaattc	tgttttatca	2760
gaccgcttct	gcgttctgat	ttaatctgta	tcaggctgaa	aatcttctct	catccgcca	2820
aacagaagct	agcggccgat	ccccaaaaa	aaaaaaaaa	aaaaaaaaa	gagtcagag	2880
tggccccgcc	gttcgcgcgc	gggggggggg	gggggggggg	acactttcgg	acatctggtc	2940
gacctccagc	atcgggggaa	aaaaaaaaa	caaagtctcg	cccggagtag	tggtcgacct	3000
ccgaagttgg	gggggagtag	aaacagggta	gataatcact	cactgacgta	cgttgagcaa	3060
ctgactgaaa	tgccttgagc	aactgactga	aatgcctgac	gtcttttagca	aaagcaggg	3120
agataatcac	tactgagtg	acatccacat	cgtaccagga	ttggctgaga	cgaaaaacat	3180
attgtaccag	ggattggctg	agacgaaaaa	catattgtag	gtacccaaat	gaacactcaa	3240
atcctggttt	tcgcccttgc	ggcagtcac	cccacaaatg	cagacaaat	ttgtcttgga	3300
catcatgctg	tatcaaatgg	caccaaagta	aacacactca	ctgagagagg	agtagaagtt	3360
gtcaatgcaa	cggaaacagt	ggagcggaca	aacatcccca	aaatttgctc	aaaagggaaa	3420
agaaccactg	atcttgacca	atgcggactg	ttagggacca	ttaccggacc	acctcaatgc	3480
gaccaatttc	tagaattttc	agctgatcta	ataatcgaga	gacgagaagg	aatgatgtt	3540
tgttaccggg	ggaagtttgt	taatgaagag	gcattgcgac	aaatcctcag	aggatcaggt	3600
gggattgaca	aagaaacaat	gggattcaca	tatagtggaa	taaggacca	cggaaacact	3660
agtgcagtga	gaagatcagg	gtcttcattc	tatgcagaaa	tggagtggct	cctgtcaaat	3720
acagacaatg	cttctttccc	acaaatgaca	aaatcataca	aaaacacagg	gagagaatca	3780
gctctgatag	tctggggaat	ccaccattca	ggatcaacca	ccgaacagac	caaactatat	3840
gggagtggaa	ataaactgat	aacagtcggg	agttccaaat	atcatcaatc	ttttgtgccg	3900
agtccaggaa	cacgaccgca	gataaatggc	cggtcgggac	ggattgattt	tcattgggtg	3960
atcttggaac	ccaatgatac	agttactttt	agtttcaatg	gggctttcat	agctccaaat	4020
cgtgccagct	tcttgagggg	aaagtccatg	gggatccaga	gcgatgtgca	ggttgatgct	4080
aattgcgaag	gggaatgcta	ccacagtgga	gggactataa	caagcagatt	gccttttcaa	4140
aacataaata	gcagagcagt	tggcaaatgc	ccaagatatg	taaaacagga	aagtttatta	4200
ttggcaactg	ggatgaagaa	cgttcccga	ccttccaaaa	aaaggaaaaa	aagaggcctg	4260
tttggtgcta	tagcaggggt	tattgaaaat	ggttgggaa	gtctggctga	cgggtggtac	4320
ggtttcaggc	atcagaatgc	acaaggagaa	ggaactgcag	cagactacaa	aagcacccaa	4380
tcggcaattg	atcagataac	cggaaagtta	aatagactca	ttaagaaaac	caaccagcaa	4440
tttgagctaa	tagataatga	attcactgaa	gtggaaaagc	agattggcaa	tttaattaac	4500
tggaccaaag	actccatcac	agaagtatgg	tcttacaatg	ctgaacttct	tgtggcaatg	4560
gaaaaccagc	acactattga	tttggctgat	tcagagatga	acaagctgta	tgagcagagt	4620
aggaaacaat	taagggaata	tgctgaagag	gatggcactg	gttgctttga	aatttttcat	4680
aaatgtgacg	atgattgtat	ggctagtata	aggaacaata	cttatgatca	cagcaaatac	4740
agagaagaag	cgatgcaaaa	tagaatacaa	attgaccag	tcaaattgag	tagtggctac	4800
aaagatgtga	tactttggtt	tagcttcggg	gcatcatgct	ttttgcttct	tgccattgca	4860
atgggccttg	ttttcatatg	tgtgaagaac	ggaaacatgc	ggtgcactat	atgcatttaa	4920
agcttgcatg						4930

&lt;210&gt; 2

&lt;211&gt; 5811

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 2

aattcctttg	cctaatttaa	atgaggactt	aacctgtgga	aatattttga	tgtgggaagc	60
tgttactgtt	aaaactgagg	ttattggggt	aactgctatg	ttaaacttgc	attcagggac	120
acaaaaaact	catgaaaatg	gtgctggaaa	acccattcaa	gggtcaaatt	ttcatttttt	180
tgctgttggt	ggggaacctt	tggagctgca	gggtgtgtta	gcaaactaca	ggaccaaata	240
tcctgctcaa	actgtaaccc	caaaaaatgc	tacagttgac	agtcagcaga	tgaacactga	300
ccacaaggct	gttttgata	aggataatgc	ttatccagtg	gagtgtggg	ttcctgatcc	360
aagtaaaaat	gaaaacacta	gatatttttg	aacctacaca	gggtgggaaa	atgtgcctcc	420
tgttttgcac	attactaaca	cagcaaccac	agtgtcttct	gatgagcagg	gtgttgggcc	480
cttggtgcaa	gctgacagct	tgtatgtttc	tgctgttgac	atttgtgggc	tgtttaccac	540
cacttctgga	acacagcagt	ggaagggact	tcccagatat	tttaaaatta	cccttagaaa	600
gcggtctgtg	aaaaacccct	aaaaaatttc	ctttttgtta	agtgcctaa	ttaacaggag	660
gacacagagg	gtggatgggc	agcctatgat	tggaatgtcc	tctcaagtag	aggaggttag	720
ggtttatgag	gacacagagg	agcttcctgg	ggatccagac	atgataagat	acattgatga	780
gtttgacaa	accacaacta	gaatgcagtg	aaaaaaatgc	tttattttgtg	aaatttgtga	840
tgctattgct	ttatttgtaa	ccattataag	ctgcaataaa	caagttaaca	acaacaattg	900
cattcatttt	atgtttcagg	ttcaggggga	ggtgtgggag	gtttttttaa	gcaagtaaaa	960
cctctacaaa	tgtggatgag	ctgattatga	tctctagtca	aggcactata	catcaaatat	1020
tccttattaa	cccctttaca	aattaaaaag	ctaaaggtac	acaatttttg	agcatagtta	1080
ttaatagcag	acactctatg	cctgtgtgga	gtaagaaaaa	acagtatgtt	atgattataa	1140
ctgttatgcc	tacttataaa	ggttacagaa	tatttttcca	taattttctt	gtatagcagt	1200
gcagcttttt	cctttgtggt	gtaaatagca	aagcaagcaa	gagttctatt	actaaacaca	1260
gcatgactca	aaaaacttag	caattctgaa	ggaaagtcct	tggggctctc	tacctttctc	1320
ttcttttttg	gaggagtaga	atgttgagag	tcagcagtag	cctcatcatc	actagatggc	1380
atttcttctg	agcaaaacag	gttttcctca	ttaaaggcat	tccaccactg	ctccattcca	1440
tcagttccat	aggttggaa	ctaaaataca	caaacaatta	gaatcagtag	tttaacacat	1500
tatacactta	aaaattttat	atttacctta	gagctttaaa	tctctgtagg	tagtttgtcc	1560
aattatgtca	caccacagaa	gtaaggttcc	ttcacaaaga	tccgggacca	aagcggccat	1620
cgtgcctccc	cactcctgca	gttcgggggc	atggatgcgc	ggatagccgc	tgctggtttc	1680
ctggatgccg	acggatttgc	actgccggta	gaactccgcg	aggtcgtcca	gcctcaggca	1740
gcagctgaac	caactcgcga	ggggatcgag	cccggggtgg	gcgaagaact	ccagcatgag	1800
atccccgcgc	tggaggatca	tccagccggc	gtcccggaaa	acgattccga	agcccaacct	1860
ttcatagaag	gcggcggtgg	aatcgaaatc	tcgtgatggc	aggttgggcg	tcgcttggtc	1920
ggtcatttctg	atgaattcga	gtcgggtacc	cggggatcct	ctagaggcat	ttcagtttctg	1980
tcctcacgga	ctcatcagag	ttgtcgaatt	cgaacccag	agtcctcgtc	agaagaactc	2040
gtcaagaagg	cgatagaagg	cgatgcgctg	cgaatcggga	gcggcgatac	cgtaaagcac	2100
gaggaagcgg	tcagcccat	cgccgccaag	ctcttcagca	atatcacggg	tagccaacgc	2160
tatgtcctga	tagcgggtccg	ccacacccag	ccggccacag	tcgatgaatc	cagaaaagcg	2220
gccattttcc	accatgatat	tcggcaagca	ggcatcgcca	tgggtcacga	cgagatcctc	2280
gccgtcgggc	atgcgcgcct	tgagcctggc	gaacagttcg	gctggcgcg	gcccctgatg	2340
ctcttcgtcc	agatcatcct	gatcgacaag	accggcttcc	atccgagtag	gtgtcgtctc	2400
gatgcgatgt	ttcgcttggt	ggtcgaatgg	gcaggtagcc	ggatcaagcg	tatgcagccg	2460
ccgcattgca	tcagccatga	tggatacttt	ctcggcagga	gcaaggtgag	atgacaggag	2520
atcctgcccc	ggcacttcgc	ccaatagcag	ccagtcctct	cccgttcag	tgacaacgtc	2580
gagcacagct	gcgcaaggaa	cgcccgctcg	ggccagccac	gatagccgcg	ctgcctcgtc	2640
ctgcagttca	ttcagggcac	cggacaggtc	ggtcttgaca	aaaagaaccg	ggcgcccctg	2700
cgctgacagc	cgggaacacg	cggcatcaga	gcagccgatt	gtctgttgtg	cccagtcata	2760
gccgaatagc	ctctccaccc	aagcggccgg	agaacctgcg	tgcaatccat	cttgttcaat	2820
catgcgaaac	gatactcatc	ctgtctcttg	atcagatctt	gatccccctg	gccatcagat	2880
ccttgggcggc	aagaaagcca	tccagtttac	tttgacgggc	ttcccaacct	taccagaggg	2940
cgccccagct	ggcaattccg	gttcgcttgc	tgtccataaa	accgcccag	ctagctatcg	3000
ccatgtaagc	ccactgcaag	ctacctgctt	tctcttttgcg	cttgcgtttt	cccttgctcca	3060
gatagcccag	tagctgacat	tcacccgggg	tcagcacctg	ttctgcggac	tggctttcta	3120
cgtgttccgc	ttccttttagc	agcccttgcg	ccctgagtag	ttgcggcagc	gtgaagcttt	3180

29

```

ttgcaaaagc ctaggcctcc aaaaaagcct cctcactact tctggaatag ctcagaggcc 3240
gaggcggcct cggcctctgc ataaataaaa aaaattagtc agccatgggg cggagaatgg 3300
gcggaactgg gcggagttag gggcgggatg ggcggagtta ggggcgggac tatggttgct 3360
gactaattga gatgcatgct ttgcatactt ctgcctgctg gggagcctgg ggactttcca 3420
cacctggttg ctgactaatt gagatgcatg ctttgcatac ttctgcctgc tggggagcct 3480
ggggactttc cacaccctaa ctgacacaca ttccacagct gcctcgcgcg ttctcgggtgat 3540
gacggtgaaa acctctgaca catgcagctc ccggagacgg tcacagcttg tctgtaagcg 3600
gatgccggga gcagacaagc ccgtcagggc gcgtcagcgg gtgttgccgg gtgtcggggc 3660
gcagccatga cccagtcacg tagcgatagc ggagtgtata ctggcttaac tatgcggcat 3720
cagagcagat tgtactgaga gtgcaccata tgcggtgtga aataccgcac agatgcgtaa 3780
ggagaaaata ccgcatcagg cgctcttccg cttcctcgct cactgactcg ctgcgctcgg 3840
tcgttcggct gcggcgagcg gtatcagctc actcaaaggc ggtaatacgg ttatccacag 3900
aattcaggga taacgaagga aagaacatgt gagcaaaagg ccagcaaaag gccaggaacc 3960
gtaaaaaggc cgcgttgctg gcgtttttcc ataggtcccg cccccctgac gagcatcaca 4020
aaaatcgacg ctcaagtcag aggtggcgaa acccgacagg actataaaga taccaggcgt 4080
ttccccctgg aagctccctc gtgcgctctc ctgttccgac cctgccgctt accggatacc 4140
tgtecgccctt tctcccttcg ggaagcgtgg cgctttctca tagctcacgc tgtaggtatc 4200
tcagttcggg gttaggtcgtt cgctccaagc tgggctgtgt gcacgaacc cccgttcagc 4260
ccgaccgctg cgccttatcc ggtaactatc gtcttgagtc caacccggtg agacacgact 4320
tatcgccact ggcagcagcc actggtaaca ggattagcag agcgaggatg gtaggcgggtg 4380
ctacagagtt cttgaagtgg tggcctaact acggctacac tagaaggaca gtatttggtg 4440
tctgcgctct gctgaagcca gttaccttcg gaaaaagagt tggtagctct tgatccggca 4500
aacaaccac cgtcggtagc ggtggttttt ttgtttgcaa gcagcagatt acgcgcagaa 4560
aaaaaggatc tcaagaagat cctttgatct tttctacggg gtctgacgct cagtggaaacg 4620
aaaactcacg ttaagggatt ttggtcatga gattatcaaa aaggatcttc acctagatcc 4680
ttttaaatta aaaatgaagt tttaaatcaa tctaaagtat atatgagtaa acttgggtctg 4740
acagttacca atgcttaatc agtgaggcac ctatctcagc gatctgtcta ttctgttcat 4800
ccatagttgc ctgactcccc gtcgtgtaga taactacgat acgggagggc ttaccatctg 4860
gccccagtg cgaatgata ccgcgagacc cacgctcacc ggctccagat ttatcagcaa 4920
taaaccagcc agccggaagg gccgagcgca gaagtgggtc tgcaacttta tccgcctcca 4980
tccagtctat taattggttc cgggaagcta gagtaagtag ttcgccagtt aatagtttg 5040
gcaacgttgt tgccattgct gcaggcatcg tgggtgtcag ctcgtcgttt ggtatggctt 5100
cattcagctc cggttcccaa cgatcaaggc gagttacatg atcccccatg ttgtgcaaaa 5160
aagcggttag ctcttcgggt cctccgatcg ttgtcagaag taagttggcc gcagtgttat 5220
cactcatggt tatggcagca ctgcataatt ctcttactgt catgccatcc gtaagatgct 5280
tttctgtgac tggtagtac tcaaccaagt cattctgaga atagtgtatg cggcgaccga 5340
gttgctcttg cccggcgta acacgggata ataccgcgcc acatagcaga actttaaaag 5400
tgctcatcat tggaaaacgt tcttcggggc gaaaactctc aaggatctta ccgctgttga 5460
gatccagttc gatgtaacc actcgtgcac ccaactgatc ttcagcatct tttactttca 5520
ccagcgtttc tgggtgagca aaaacaggaa ggcaaaatgc cgcaaaaaag ggaataaggg 5580
cgacacggaa atgttgaaata ctcatactct tcctttttca atattattga agcatttatc 5640
agggttattg tctcatgagc ggatacatat ttgaatgtat ttagaaaaat aaacaaatag 5700
gggttcgcg cactttccc cgaaaagtgc cacctgacgt ctaagaaacc attattatca 5760
tgacattaac ctataaaaat aggcgtatca cgaggccctt tcgtcttcaa g 5811

```

&lt;210&gt; 3

&lt;211&gt; 2005

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 3

```

agtagaaaca agggatattt tctttaccta gctaggcctg cgcgcaatta accctcacta 60
aagggaacaa aagctggagc tccaccgcgg tggcggccgc tctagcatat gtttttcgtc 120
tcagccaatc cctctagcat atgtttttcg tctcagccaa tccctctaga agcttcgtac 180
gcatgcaagc tttaaatgca tatagtgcac cgcatgttcc cgttcttcac acatatgaaa 240

```



30

```

acaaggccca ttgcaatggc aagaagcaaa aagcatgatg ccccgaaagt aaaccaaaagt 300
atcacatctt tgtagccact actcaatttg actgggtcaa tttgtattct attttgcac 360
gcttcttctc tgtatttgct gtgatcataa gtattgttcc ttatactagc catacaatca 420
tcgtcacatt tatgaaaaat ttcaaagcaa ccagtgccat cctcttcagc attttccctt 480
aattgtttcc tcaactcgctc atacagcttg ttcactctcg aatcagccaa atcaatagt 540
tgctggtttt ccattgccac aagaagttca gcattgtaag accatacttc tgtgatggag 600
tctttgggtcc agttaattaa attgccaatc tgcttttcca cttcagtga ttcattatct 660
attagctcaa attgctgggt ggttttctta atgagtctat ttaactttcc gggtatctga 720
tcaattgccg attgggtgct tttgtagtct gctgcagttc cttctccttg tgcattctga 780
tgctgaaac cgtaccacc gtcgaccaga cttcccaac ctttttcaat aaacctgct 840
atagcaccia acaggcctct ttttttccct tttttggaag gttcgggaac gttcttcac 900
ccagttgcca ataataaact ttctgtttt acatatcttg ggcatttgcc aactgctctg 960
ctatttatgt tttgaaaagg caatctgctt gttatagtaa ctcaactgtg gtagcattcc 1020
ccttcgcaat tagcatcaac ctgcacatcg ctctggatcc ccattggactt tccctcaag 1080
aagctggcac gatttgagc tatgaaagcc ccattgaaac taaaagtaac tgtatcattg 1140
ggatccaaga tcaaccaatg aaaatcaatc cgtccggacc ggccatttat ctgcggctgt 1200
gttctggac tcggcacaaa agattgatga tatttggaac tcccgaactgt tatcagttta 1260
tttccactcc catatagttt ggtctgttcg gtggttgatc ctgaatgggtg gattccccag 1320
actatcagag ctgattctct cctgtgttt ttgtatgatt ttgtcatttg tgggaaagaa 1380
gcattgtctg tatttgacag gagccactcc atttctgcat agaatagaaga ccctgatctt 1440
ctacatgcac tagttgttcc gttggtcctt attccactat atgtgaatcc cattgtttct 1500
ttgtcaatcc cacctgatcc tctgaggatt tgcgcaatg cctcttcatt aacaaacttc 1560
cccgggtaac aaacatcatt tcttctcgt ctctcgatta ttagatcagc tgaaaattct 1620
agaaattggt cgcattgagg tggtcggta atggtcctta acagtcgca ttggccaaga 1680
tcagtgggtc ttttcccttt tgagcaaatt ttggggatgt ttgtccgctc cactgtttcc 1740
gttgcatga caacttctac tctctctca gtgagtgtgt ttactttggt gccatttgat 1800
acagcatgat gtccaagaca aattttgtct gcatttgtgg ggatgactgc cgcaaggcg 1860
aaaaccagga tttgagtgtt cattttggta cctacaatat gttttctgct tcagccaatc 1920
cctggtacaa tatgttttct gtctcagcca atcctggtac gatgtggatg tcaactcagt 1980
agtgattatc taccctgctt ttgct 2005

```

&lt;210&gt; 4

&lt;211&gt; 1146

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 4

```

agtagaaaca agggatattt tctttaccta gctaggcctg cgcgcaatta accctcacta 60
aagggaacaa aagctggagc tccaccgagg tggcgccgc tctagcatat gttttctgct 120
tcagccaatc cctctagcat atgtttttcg tctcagccaa tccctctaga agcttcgtac 180
gcatgcttaa ataagctgaa acgagaaagt tcttatctct tgctccactt caagcggtag 240
ttgtaaggct tgcataaatg ttatttggtc aaaactatct tctgttatct tcaatctatg 300
tctcacttct tcaattaacc atcttatttc ttcaaatttc tgactcaatt gttctcgcca 360
ttttccgttt ctgctttgga gggagtggag gtcccccatt ctactactg cttctccaag 420
cgaatctctg tatagtttca gagactcgaa ctgtgttate attccattca agtccctcca 480
tgaggacccc aattgcattt ttgacatcct catcagtagt tccctggaaga gaaggcaatg 540
gtgaaatttc gccgacaatt gctccctcat cgggttaaagc ccttaatagt atgagagttt 600
ccagccgatc gaaaatcaca ctgaagtttg ctttcagtat gatgttcttc cccatgatcg 660
cctgggtccat tctgatgcaa agggagcctg ccactttctg tttgggcatg agcatgaacc 720
agtcccttga catctcttca agagtcatgt cagttaggta gcgtgtagca ggtacagagg 780
caatggctcat ttttaagtgc tcatcggtt cgtcctccag aatccgctcc actatctgct 840
ttccaacacg agtagctgtg tcgatgtcca gaccaagagt gctgcctctt cccctcaggg 900
acttctgate tcggcgaagt cgggtcaagga atggggcact acccatttct tgggtctgcaa 960
atcgtttgct gacatgccaa agaaagcagt ctacctgaaa gcttgacaca gtgttggaat 1020
ccattatggt acctacaata tgtttttcgt ctcagccaat cctggttaca atatgttttt 1080

```



31

cgtctcagcc aatcctggta c gatgtggat gtcactcagt gactgattat ctaccctgct 1140  
tttgct 1146

&lt;210&gt; 5

&lt;211&gt; 5860

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 5

catcgattgg ctgactgatg agtccgtgag gacgaaacga aaaacatatt gtagagctcg 60  
aattcatcga aatgaccgac caagcgacgc ccaacctgcc atcacgagat ttcgattcca 120  
ccgcccgcctt ctatgaaagg ttgggcttcg gaatcgtttt ccgggacgcc ggctggatga 180  
~~tctccagcg aggggatctg atgctggagt tcttcgccc cccggggctc gatccctcg 240~~  
cgagttgggtt cagctgctgc ctgaggctgg acgacctgcg ggagttctac cggcagtgca 300  
aatccgtcgg catccaggaa accagcagcg gctatccgcg catccatgcc cccgaactgc 360  
aggagtgggg aggacgatg gccgctttgg tcccgatct ttgtgaagga accttacttc 420  
tgtggtgtga cataattgga caaactacct acagagattt aaagctctaa ggtaaatata 480  
aaatttttaa gtgtataatg tggttaaacta ctgattctaa ttgtttgtgt attttagatt 540  
ccaacctatg gaactgatga atgggagcag tgggtggaatg cctttaatga ggaaaacctg 600  
ttttgctcag aagaaatgcc atctagtgat gatgaggcta ctgctgactc tcaacattct 660  
actcctccaa aaaagaagag aaaggtagaa gaccccaagg actttccttc agaattgcta 720  
agttttttga gtcatgctgt gtttagtaat agaactcttg cttgctttgc tatttacacc 780  
acaaaggaaa agctgcact gctatacaag aaaattatgg aaaaatattc tgtaaccttt 840  
ataagtaggc ataacagtta taatcataac atactgtttt ttcttactcc acacaggcat 900  
agagtgtctg ctattaataa ctatgctcaa aaattgtgta cctttagctt tttaatttgt 960  
aaaggggtta ataaggaata tttgatgtat agtgccttga ctagagatca taatcagcca 1020  
taccacattt gtagagggtt tacttgcttt aaaaaacctc ccacacctcc cctgaacct 1080  
gaaacataaa atgaatgcaa ttgttgttgt taacttgttt attgcagctt ataatgggtta 1140  
caaataaagc aatagcatca caaatttcac aaataaagca tttttttcac tgcattctag 1200  
ttgtggtttg tccaaactca tcaatgtatc ttatcatgtc tggatcccca ggaagctcct 1260  
ctgtgtcctc ataaacccta acctcctcta cttgagagga cattccaatc ataggctgcc 1320  
catccacct ctgtgtcctc ctgttaatta ggtcacttaa caaaaaggaa attgggtagg 1380  
ggtttttcac agaccgcttt ctaagggtaa ttttaaaata tctgggaagt ccctccact 1440  
gctgtgttcc agaagtgttg gtaaacagcc cacaaatgtc aacagcagaa acatacaagc 1500  
tgtcagcttt gcacaagggc ccaacacct gctcatcaag aagcactgtg gttgctgtgt 1560  
tagtaatgtg caaaacagga ggcacatttt cccacctgt gtaggttcca aaatatctag 1620  
tgttttcatt tttacttgga tcaggaacct agcactccac tggataagca ttatccttat 1680  
ccaaaacagc cttgtggtca gtgttcact gctgactgtc aactgtagca ttttttgggg 1740  
ttacagtttg agcaggatat ttggctcctgt agtttgctaa cacacctgc agctccaaag 1800  
gttccccacc aacagcaaaa aaatgaaaat ttgaccttg aatgggtttt ccagcaccat 1860  
tttcatgagt tttttgtgtc cctgaatgca agtttaacat agcagttacc ccaataacct 1920  
cagttttaac agtaacagct tcccacatca aaatatttcc acaggttaag tctcattta 1980  
aattaggcaa aggaattctt gaagacgaaa gggcctcgtg atacgctat ttttataggt 2040  
taatgtcatg ataataatgg tttcttagac gtcagggtggc acttttcggg gaaatgtgcg 2100  
cggaaccctt atttgtttat ttttctaaat acattcaaat atgtatccgc tcatgagaca 2160  
ataacctga taaatgcttc aataatattg aaaaaggag agtatgagta ttcaacattt 2220  
ccgtgtcgcc cttatccct tttttgcggc attttgcctt cctgtttttg ctacccaga 2280  
aacgctggtg aaagtaaaag atgctgaaga tcagttgggt gcacgagtgg gttacatcga 2340  
actggatctc aacagcggtg agatccttga gagttttcgc cccgaagaac gttttccaat 2400  
gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccggtgtg acgcccggca 2460  
agagcaactc ggtcgccgca tacactattc tcagaatgac ttggttgagt actcaccagt 2520  
cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg ctgccataac 2580  
catgagtgat aacactgcgg ccaacttact tctgacaacg atcggaggac cgaaggagct 2640  
aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcgtt gggaaccgga 2700  
gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgcctgcag caatggcaac 2760

aacgttgccg aaactattaa ctggcgaact acttactcta gcttcccggc aacaattaat 2820  
agactggatg gaggcggata aagttgcagg accacttctg cgctcggccc ttccggctgg 2880  
ctggtttatt gctgataaat ctggagccgg tgagcgtggg tctcgcggtt tcattgcagc 2940  
actggggcca gatggtaagc cctcccgtat cgtagttatc tacacgacgg ggagtcaggc 3000  
aactatggat gaacgaaata gacagatcgc tgagataggt gcctcactga ttaagcattg 3060  
gtaactgtca gaccaagttt actcatatat acttttagatt gatttaaaac ttcatTTTTA 3120  
atttaaaagg atctaggtga agatcctttt tgataatctc atgacaaaaa tcccttaacg 3180  
tgagtttttcg ttccactgag cgtcagaccc cgtagaaaag atcaaaggat cttcttgaga 3240  
tccttttttt ctgcgcgtaa tctgctgctt gcaaacaaaa aaaccaccgc taccagcggg 3300  
ggtttggttg ccggatcaag agctaccaac tctttttccg aaggtaactg gcttcagcag 3360  
agcgcagata ccaaatactg tcctttctagt gtagccgtag ttaggccacc acttcaagaa 3420  
ctctgtagca ccgcctacat acctcgtctt gctaatectg ttaccagtgg ctgctgccag 3480  
~~tgccgataag tegtgtctta cccggttggg ctaagaaga tagttaccgg ataaggcgca 3540~~  
gcggtcgggc tgaacggggg gtctgtgcac acagcccagc ttggagcgaa cgacctacac 3600  
cgaactgaga tacctacagc gtgagctatg agaaagcgcc acgcttcccg aaggagaaaa 3660  
ggcggacagg tatccggtaa gcggcagggg cggaacagga gagcgcacga gggagcttcc 3720  
agggggaaac gcctgggtatc tttatagtcg tgctcgggtt cgccacctct gacttgagcg 3780  
tcgatttttg tgatgctcgt caggggggcg gagcctatgg aaaaacgcca gcaacgcggc 3840  
ctttttacgg ttcttggect tttgctggcc ttttgctcac atgttcttct ctgcgttatc 3900  
ccctgattct gtggataacc gtattaccgc ctttgagtga gctgataccg ctgcgcgcag 3960  
ccgaacgacc gagcgcagcg agtcagttag cgaggaagcg gaagagcgcc tgatgcggta 4020  
ttttctcctt acgcatctgt gcggtatttc acaccgata tgggtgactc tcagtacaat 4080  
ctgctctgat gccgcatagt taagccagta tacactccgc tatcgctacg tgactgggtc 4140  
atggctgcgc cccgacaccc gccaacaccc gctgacgcgc cctgacgggc ttgtctgctc 4200  
ccggcatccg cttacagaca agctgtgacc gtctccggga gctgcatgtg tcagaggttt 4260  
tcaccgtcat caccgaaacg cgcgaggcag ctgtggaatg tgtgtcagtt aggggtgtgga 4320  
aagtccccag gctccccagc aggcagaagt atgcaaagca tgcattctca ttagtcagca 4380  
accagggtgtg gaaagtcccc aggtccccca gcaggcagaa gtatgcaaag catgcattct 4440  
aattagtcag caaccatagt cccgccccta actccgccc tccgcccct aactccgcc 4500  
agttccgccc attctccgcc ccatggctga ctaatttttt ttatttatgc agaggccgag 4560  
gccgcctcgg cctctgagct attccagaag tagtgaggag gcttttttgg aggcctaggc 4620  
ttttgcaaaa agcttcacgc tgccgcaagc actcagggcg caagggtgc taaaggaagc 4680  
ggaacacgta gaaagccagt ccgcagaaac ggtgctgacc ccggatgaat gtcagctact 4740  
gggctatctg gacaagggaa aacgcaagcg caaagagaaa gcaggtagct tgcagtgggc 4800  
ttacatggcg atagctagac tgggcgggtt tatggacagc aagcgaaccg gaattgccag 4860  
ctggggcgcc ctctggtaag gttgggaagc cctgcaaagt aaactggatg gctttcttgc 4920  
cgccaaggat ctgatggcg aggggatcaa gatctgatca agagacagga tgaggatcgt 4980  
ttcgcattgat tgaacaagat ggattgcacg caggttctcc ggccgcttgg gtggagagcg 5040  
tattcggcta tgactggga caacagacaa tcggctgctc tgatgccgcc gtgttccggc 5100  
tgtcagcgca ggggcgccc gttctttttg tcaagaccga cctgtccggt gccctgaatg 5160  
aactgcagga cgaggcagcg cggctatcgt ggctggccac gacgggctt ccttgccgag 5220  
ctgtgctcga cgttgtcact gaagcgggaa gggactggct gctattgggc gaagtgccgg 5280  
ggcaggatct cctgtcatct caccttgctc ctgccgagaa agtatccatc atggctgatg 5340  
caatgcggcg gctgcatacg cttgatccgg ctacctgcc attcgaccac caagcgaaac 5400  
atcgcatcga gcgagcacgt actcggatgg aagccggtct tgctgatcag gatgatctgg 5460  
acgaagagca tcaggggctc gcgccagccg aactgttcgc caggctcaag gcgcgcagtc 5520  
ccgaecggca ggatctcgtc gtgacctatg gcgatgctg cttgccgaat atcatggtgg 5580  
aaaatggccg cttttctgga ttcacgact gtggccggct ggggtgtggc gaccgctatc 5640  
aggacatagc gttggctacc cgtgatattg ctgaagagct tggcggcgaa tgggctgacc 5700  
gcttctcgt gctttacgg atcgccgctc ccgattcgca gcgcacgcc ttctatcgcc 5760  
ttcttgacga gttcttctga gcgggactct ggggttcgaa tcctaccagg gattggetga 5820  
ctgatgagtc cgtgaggacg aaacgaaaaa catatggtac 5860

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 6

```
gaggcatttc agtcagttgc tcaaggtacc aaaatgaaca ctcaaactct ggtttttcgcc 60
cttgcggcag tcacccccac aaatgcagac aaaatttgtc ttggacatca tgctgtatca 120
aatggcacca aagtaaacac actcactgag agaggagtag aagttgtcaa tgcaacggaa 180
acagtggagc ggacaaacat ccccaaaatt tgctcaaaag ggaaaagaac cactgatctt 240
ggccaatgcg gactgttagg gaccattacc ggaccacctc aatgcgacca atttctagaa 300
ttttcagctg atctaataat cgagagacga gaaggaaatg atgtttgtta cccgggggaag 360
tttgttaatg aagaggcatt gcgacaaatc ctcagaggat caggtgggat tgacaaagaa 420
acaatgggat tcacatatag tggaataagg accaacggaa caactagtgc atgtagaaga 480
tcagggtctt cattctatgc agaaatggag tggctcctgt caaatacaga caatgcttct 540

ttcccacaaa tgacaaaatc atacaaaaac acaggggagag aatcagctct gatagtctgg 600
ggaatccacc attcaggatc aaccaccgaa cagaccaaac tatatgggag tggaaataaa 660
ctgataacag tcgggagttc caaatatcat caatcttttg tgccgagtc aggaacacga 720
ccgcagataa atggccggtc cggacggatt gattttcatt ggttgatctt ggatcccaat 780
gatacagtta ctttttagtt caatggggct tcatagctc caaatcgtgc cagcttcttg 840
aggggaaagt ccatggggat ccagagcgat gtgcaggttg atgctaattg cgaaggggaa 900
tgctaccaca gtggagggac tataacaagc agattgcctt ttcaaacat aaatagcaga 960
gcagttggca aatgcccaag atatgtaaaa caggaaagt tattattggc aactgggatg 1020
aagaacgttc ccgaacctc caaaaaaagg aaaaaaagag gctgttttg tgctatagca 1080
gggtttattg aaaatggttg ggaaggtctg gtcgacgggt ggtacggttt caggcatcag 1140
aatgcacaag gagaaggaa tgcagcagac taaaaaagca cccaatcggc aattgatcag 1200
ataaccggaa agttaaatag actcattaag aaaaccaacc agcaatttga gctaatagat 1260
aatgaattca ctgaagtgga aaagcagatt ggcaatttaa ttaactggac caaagactcc 1320
atcacagaag tatggtctta caatgctgaa cttcttgttg caatggaaaa ccagcacact 1380
attgatttgg ctgattcaga gatgaacaag ctgtatgagc gagtgaggaa acaattaagg 1440
gaaaatgctg aagaggatgg cactgggtgc tttgaaattt ttcataaatg tgacgatgat 1500
tgtatggcta gtataaggaa caatacttat gatcacagca aatacagaga agaagcgatg 1560
caaaatagaa tacaaattga cccagtcaaa ttgagtatg gctacaaaga tgtgatactt 1620
tggttttagct tcggggcatc atgctttttg cttcttgcca ttgcaatggg ccttgttttc 1680
atatgtgtga agaacggaaa catgcgggtc actatttgta tataggtttg gaaaaaaca 1740
ccccttgttt ctactcccc ccaacttcgg aggtcgacca gtactcggg cgaaactttg 1800
tttttttttt ttccccgat gctggagggtc gaccagatgt ccgaaagtgt ccccccccc 1860
cccccccccc ggcgcggaac ggcggggcca ctctggactc tttttttttt tttttttttt 1920
ttttttgggg atcgccgct agcttctgtt ttggcggtg agagaagatt ttcagcctga 1980
tacagattaa atcagaacgc agaagcggtc tgataaaaca gaatttgcct ggcggcagta 2040
gcgcgggtgt cccacctgac cccatgccga actcagaagt gaaacgccgt agcgccgatg 2100
gtagtgtggg gtctcccat gcgagagtag ggaactgcca ggcacaaat aaaacgaaag 2160
gctcagtcga aagactgggc ctttcgtttt atctgttgtt tgcgggtgaa cgctctcctg 2220
agtaggacaa atccgcggg agcggatttg aacgttgcca agcaacggcc cggagggtgg 2280
cgggcaggac gcccgccata aactgccagg catcaaatta agcagaaggc catcctgacg 2340
gatggccttt ttgcgtttct acaaactctt ttgtttattt ttctaaatac attcaaatat 2400
gtatccgctc atgagacaat aaccctgata aatgcttcaa taatattgaa aaaggaagag 2460
tatgagtatt caacatttcc gtgtcgccct tattcccttt tttgcggcat tttgccttcc 2520
tgtttttgct caccagaaa cgctggtgaa agtaaaagat gctgaagatc agttgggtgc 2580
acgagtgggt tacatcgaa tggatctcaa cagcggtaag atccttgaga gttttcgccc 2640
cgaagaacgt tttccaatga tgagcacttt taaagttctg ctatgtggcg cggattatc 2700
ccgtgttgac gccgggcaag agcaactcgg tcgccgcata cactattctc agaatgactt 2760
ggttgagtac tcaccagtca cagaaaagca tcttacggat ggcacgacag taagagaatt 2820
atgcagtgtc gccataacca tgagtataa cactgcggcc aacttacttc tgacaacgat 2880
cggaggaccg aaggagctaa ccgctttttt gcacaacatg ggggatcatg taactcgcc 2940
tgatcgttgg gaaccggagc tgaatgaagc cataccaaac gacgagcgtg acaccacgat 3000
gcctgtagca atggcaacaa cgttgcgcaa actattaact ggcgaactac ttactctagc 3060
```

34

```
ttccccggcaa caattaatag actggatgga ggcggataaa gttgcaggac cacttctgcg 3120
ctcgccctt cggctggct ggtttattgc tgataaatct ggagccgggtg agcgtgggtc 3180
tcgcggtatc attgcagcac tggggccaga tggtaaagccc tcccgatcgt tagttatcta 3240
cacgacgggg agtcaggcaa ctatggatga acgaaataga cagatcgctg agataggtgc 3300
ctcactgatt aagcattggt aactgtcaga ccaagtttac tcatatatac tttagattga 3360
tttaaaactt catttttaat ttaaaaggat ctagggtgaag atcctttttg ataatctcat 3420
gacaaaatc ccttaacgtg agttttcggt ccactgagcg tcagaccccg tagaaaagat 3480
caaaggatct tcttgagatc ctttttttct gcgcgtaatc tgctgcttgc aaacaaaaaa 3540
accaccgcta ccagcgggtg tttgtttgcc ggatcaagag ctaccaactc tttttccgaa 3600
ggtaactggc ttcagcagag cgcagatacc aaatactgtc cttctagtgt agccgtagt 3660
aggccaccac ttcaagaact ctgtagcacc gcctacatac ctgcgtctgc taatcctgtt 3720
accagtggct gctgccagt gcgataagtc gtgtcttacc gggttggact caagacgata 3780
ggtaccggat aaggegeage ggtgggggtg aacgggggggt tctgtcacac agcccagctt 3840
ggagcgaacg acctacaccg aactgagata cctacagcgt gagctatgag aaagcggcac 3900
gcttcccga gggagaaagg cggacaggta tccggtaaagc ggcagggctg gaacaggaga 3960
gcgcacgagg gagcttccag ggggaaacgc ctggtatctt tatagtcctg tcgggtttcg 4020
ccacctctga cttgagcgtc gatttttgtg atgctcgtea ggggggcgga gcctatggaa 4080
aaacgccagc aacgcggcct ttttacgggt cctggccttt tgctggcctt ttgctcacat 4140
gttctttcct gcgttatccc ctgattcatt aatgcaggtc acgatccttt ctggcgagtc 4200
cccgtgcgga gtcggagagc gctccctgag cgcgtgcggc ccgagaggtc gcgcctggcc 4260
ggccttcggt cctcgtgtg tcccggtcgt aggagggggc ggcgaaaat gcttccggct 4320
cccgtctgag agacacgggc cggcccccctg cgtgtggcac gggcgcccg gagggcgctc 4380
ccggcccggc gctgctccc cgtgtgtcct ggggttgacc agagggcccc gggcgctccg 4440
tgtgtggctg cgatgggtggc gtttttgggg acaggtgtcc gtgtccgtgt cgcgcgtcgc 4500
ctgggcccgc ggcgtggtcg gtgacgcgac ctcccggccc cgggggaggt atatctttcg 4560
ctccgagtcg gcattttggg ccgcccgggt attagtagaa acaggggtac 4610
```

&lt;210&gt; 7

&lt;211&gt; 3558

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 7

```
tattagtaga aacagggtat tttttattct agtacattac gccccgccct gccactcatc 60
gcagtactgt tgtaattcat taagcattct gccgacatgg aagccatcac agacggcatg 120
atgaacctga atcgccagcg gcacagcac cttgtcgctt tgcgtataat atttgcccat 180
ggtgaaaacg ggggcgaaga agttgtccat attggccacg tttaaatcaa aactggtgaa 240
actcaccagc ggattggctg agacgaaaaa catattctca ataaaccctt tagggaaata 300
ggccagggtt tcaccgtaac acgccacatc ttgcgaatat atgtgtagaa actgccggaa 360
atcgctcgtg tattcactcc agagcgatga aaacgtttca gtttgctcat ggaaaacgg 420
gtaacaaggg tgaacactat cccatatcac cagctcaccg tctttcattg ccatacggaa 480
ttccggatga gcattcatca ggcgggcaag aatgtgaata aaggccggat aaaacttgtg 540
cttatttttc tttacggctt ttaaaaaggc cgtaatatcc agctgaacgg tctggttata 600
ggtacattga gcaactgact gaaatgcctc aaaatgttct ttacgatgcc attgggatat 660
atcaacggtg gtatatccag tgattttttt ctccatgatt atggccatta ccctgtttc 720
tactcccccc caacttcgga ggtcgaccag tactccgggc gaaactttgt tttttttttt 780
tcccccgatg ctggaggtcg accagatgtc cgaaagtgtc cccccccccc ccccccccg 840
gcgcggaacg gcggggccac tctggaactc tttttttttt tttttttttt tttttgggga 900
tcggccgcta gcttctgttt tggcggatga gagaagattt tcagcctgat acagattaaa 960
tcagaacgca gaagcggctt gataaaacag aatttgcttg gcggcagtag cgcgggtgtc 1020
ccacctgacc ccatgccgaa ctccagaagt aaacgccgta gcgccgatgg tagtgtgggg 1080
tctccccatg cgagagtagg gaactgccag gcatcaaata aaacgaaagg ctcagtcgaa 1140
agactgggccc tttcgtttta tctgttgttt gtcggtgaac gctctcctga gtaggacaaa 1200
tccgcccggga gcggatttga acgttgcgaa gcaacggccc ggaggggtggc gggcaggacg 1260
cccgccataa actgccaggc atcaaattaa gcagaaggcc atcctgacgg atggcctttt 1320
```

35

tgcgtttcta	caaactcttt	tgtttatttt	tctaaatata	ttcaaatatg	tatccgctca	1380
tgagacaata	accctgataa	atgcttcaat	aatattgaaa	aaggaagagt	atgagtattc	1440
aacattttccg	tgtcgccctt	attccctttt	ttgcggcatt	ttgccttcc	gtttttgctc	1500
acccagaaac	gctggtgaaa	gtaaaagatg	ctgaagatca	gttgggtgca	cgagtgggtt	1560
acatcgaaact	ggatctcaac	agcggtaaga	tccttgagag	ttttcgcccc	gaagaacgtt	1620
ttccaatgat	gagcactttt	aaagttctgc	tatgtggcgc	ggtattatcc	cgtgttgacg	1680
ccgggcaaga	gcaactcggg	cgccgcatac	actattctca	gaatgacttg	gttgagtact	1740
caccagtcac	agaaaagcat	cttacggatg	gcatgacagt	aagagaatta	tgcaagtgtg	1800
ccataaccat	gagtataaac	actgcggcca	acttacttct	gacaacgata	ggaggaccga	1860
aggagctaac	cgcttttttg	cacaacatgg	gggatcatgt	aactcgcctt	gatcgttggg	1920
aaccggagct	gaatgaagcc	ataccaaacg	acgagcgtga	caccacgatg	cctgtagcaa	1980
tggcaacaac	gttgcgcaaa	ctattaactg	gcgaactact	tactctagct	tcccggcaac	2040
<del>aattaataga</del>	<del>ctggatggag</del>	<del>ggggataaag</del>	<del>ttgcaggacc</del>	<del>acttctgcgc</del>	<del>tcggcccttc</del>	<del>2100</del>
cggttggtg	gtttattgct	gataaatctg	gagccggtga	gcgtgggtct	cgcggtatca	2160
ttgcagcact	ggggccagat	ggtaagccct	cccgtatcgt	agttatctac	acgacgggga	2220
gtcaggcaac	tatggatgaa	cgaaatagac	agatcgctga	gatagggtgc	tcactgatta	2280
agcattggta	actgtcagac	caagtttact	catatatact	ttagattgat	ttaaaacttc	2340
atttttaatt	taaaaggatc	taggtgaaga	tcctttttga	taatctcatg	acccaaatcc	2400
cttaacgtga	gttttcgttc	cactgagcgt	cagaccccg	agaaaagatc	aaaggatctt	2460
cttgagatcc	tttttttctg	cgcgtaatct	gctgcttgca	aacaaaaaaa	ccaccgctac	2520
cagcgggtgt	ttgtttgccg	gatcaagagc	taccaactct	ttttccgaag	gtaactggct	2580
tcagcagagc	gcagatacca	aatactgtcc	ttctagtgtg	gccgtagtta	ggccaccact	2640
tcaagaactc	tgtagcaccg	cctacatacc	tcgctctgct	aatcctgtta	ccagtggctg	2700
ctgccagtgg	cgataagtcg	tgtcttaccg	ggttggaactc	aagacgatag	ttaccggata	2760
aggcgcagcg	gtcgggctga	acgggggggt	cgtgcacaca	gcccagcttg	gagcgaacga	2820
cctacaccga	actgagatac	ctacagcgtg	agctatgaga	aagcgccacg	cttcccgaag	2880
ggagaaaggc	ggacaggat	ccggtaagcg	gcagggtcgg	aacaggagag	cgcacgaggg	2940
agcttccagg	gggaaacgcc	tggatatctt	atagtcctgt	cggttttcgc	cacctctgac	3000
ttgagcgtcg	atttttgtga	tgtctcgtcag	gggggaggag	cctatggaaa	aacgccagca	3060
acgcggcctt	tttacggttc	ctggcccttt	gctggccctt	tgctcacatg	ttctttcctg	3120
cgttatcccc	tgattcatta	atgcaggtea	cgatcccttc	tggcgagtc	ccgtgcggag	3180
tcggagagcg	ctccctgagc	gcgtgcggcc	cgagaggteg	cgccctggccg	gccttcgggtc	3240
cctcgtgtgt	cccggtcgta	ggaggggccc	gccgaaaatg	cttcgggtc	ccgctctgga	3300
gacacggggc	ggccccctgc	gtgtggcacg	ggcgccgggg	agggcgctcc	cggccccggcg	3360
ctgctcccgc	gtgtgtcctg	gggttgacca	gagggccccg	ggcgctccgt	gtgtgggtgc	3420
gatggtggcg	tttttgggga	caggtgtccg	tgtccgtgtc	gcgcgtcgcc	tgggccccgcg	3480
gcgtggtcgg	tgacgcgacc	tcccggcccc	gggggaggta	tatctttcgc	tccgagtcgg	3540
cattttgggc	cgccgggt					3558

&lt;210&gt; 8

&lt;211&gt; 4343

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 8

ctttctggcg	agtccccgtg	cggagtcgga	gagcgtccc	tgagcgcgtg	cggccccgaga	60
ggtcgcgcct	ggccggcctt	cggctccctg	tgtgtcccgg	tcgtaggagg	ggccggccga	120
aaatgcttcc	ggctcccgtc	ctggagacac	gggcccggcc	cctgcgtgtg	gcacggggcg	180
ccgggagggc	gtccccggcc	cggcgctgct	cccgcgtgtg	tcctgggggt	gaccagaggg	240
ccccggggcg	tcctgtgtgt	gctgcgatgg	tggcgttttt	ggggacagg	gtccgtgtcc	300
gtgtcgcgcg	tcgcctgggc	cggcgccgtg	gtcgggtgacg	cgacctccc	gccccggggg	360
aggtatatct	ttcgtccga	gtcggcattt	tgggccgcgg	ggttattagt	agaaacaggg	420
tattttttat	actagtaagc	tcgaaggagt	ccaccatgag	taaaggagaa	gaacttttca	480
ctggagttgt	cccaattctt	gttgaattag	atggtgatgt	taatgggcac	aaattttctg	540
tcagtggaga	gggtgaagg	gatgcaacat	acggaaaact	tacccttaaa	tttatttgca	600



ctactggaaa actacctgtt ccatggccaa cacttgtcac tactttcact tatggtgttc 660  
aatgcttttc aagataccca gatcatatga aacagcatga ctttttcaag agtgccatgc 720  
ccgaagggtta tgtacaggaa agaactatat ttttcaaaga tgacgggaac tacaagacac 780  
gtgctgaagt caagtttgaa ggtgataccc ttgttaatag aatcgagtta aaaggtattg 840  
atttttaaga agatggaaac attcttggac acaaattgga atacaactat aactcacaca 900  
atgtatacat catggctgac aagcagaaga acggaatcaa ggccaacttc aagacccgcc 960  
acaacatcga ggacggcggc gtgcagctgg ccgaccacta ccagcagaac accccaattg 1020  
gcatggccc tgccttttta ccagacaacc attacctgtc cacacaatct gccctttcga 1080  
aagatcccaa cgaaaagaga gaccacatgg tcttcttga gtttgtaaca gctgctggga 1140  
ttacacatgg catggatgaa ctatacaagg gatcccatca ccatcaccat cactaagctc 1200  
catggtctag atatctagta cattacgccc cgccctgcca ctcatcgag tactgttgta 1260  
attcattaag cattctgccc acatggaagc catcacagac ggcatgatga acctgaatcg 1320  
~~ccagcggcat cagaacttg togacttgg tataatattt gccatggtg aaaacggggg 1380~~  
cgaagaagtt gtccatattg gccacgttta aatcaaaaact ggtgaaactc acccagggat 1440  
tggcactcac aaagaacatg ttctcgatga atcctttagg gaagtaggcc aggttttcac 1500  
cgtaacacgc cacatcttgc gaatatatgt gtagaaactg ccggaaatcg tcgtggtatt 1560  
cactccagag cgatgaaaac gtttcagttt gtcctgga aacggtgtaa caagggtaga 1620  
cactatccca tatcaccagc tcaccgtctt tcattgccat acggaattcc ggatgagcat 1680  
tcacagggcg ggcaagaatg tgaataaagg ccggataaaa cttgtgctta ttttcttta 1740  
cggtctttta aaaggccgta atatccagct gaacggtctg gttataggta cattgagcaa 1800  
ctgactgaaa tgcctcaaaa tgttctttac gatgccattg ggatatatca acggtggtat 1860  
atccagtgat ttttttctcc atgattatgc aaaaaatacc cttgtttcta cccccccca 1920  
acttcggagg tcgaccagta ctccgggcga aactttgttt ttttttttcc ccccgatgct 1980  
ggaggtcgac cagatgtccg aaagtgtccc cccccccccc cccccccggc gcggaacggc 2040  
ggggccactc tggactcttt tttttttttt tttttttttt tttggggatc ggccgctagc 2100  
ttctgttttg gcggatgaga gaagattttc agcctgatac agattaaatc agaacgcaga 2160  
agcggctctga taaaacagaa tttgcctggc ggcagtagcg cgggtggtccc acctgacccc 2220  
atgccgaact cagaagtga acgccgtagc gccgatggta gtgtggggtc tccccatgcg 2280  
agagtaggga actgccaggc atcaaataaa acgaaaggct cagtcgaaag actgggcctt 2340  
tcgttttatc tgttgtttgt cggatgaacgc tctctgagt aggacaaatc cgccgggagc 2400  
ggatttgaac gttgcgaagc aacggcccgg aggggtggcg gcaggacgcc cgccataaac 2460  
tgccaggcat caaattaagc agaaggccat cctgacggat ggcctttttg cgtttctaca 2520  
aactcttttg tttatttttc taaatacatt caaatatgta tccgctcatg agacaataac 2580  
cctgataaat gtttcaataa tattgaaaaa ggaagagtat gagtattcaa catttccgtg 2640  
tcgcccttat tccctttttt gcggcatttt gccttctctg ttttgctcac ccagaaacgc 2700  
tggtgaaaagt aaaagatgct gaagatcagt tgggtgcacg agtgggttac atcgaactgg 2760  
atctcaacag cggtaagatc cttgagagtt ttcgccccga agaactgttt ccaatgatga 2820  
gcacttttaa agttctgcta tgtggcgcgg tattatcccg tgttgacgcc gggcaagagc 2880  
aactcggtcg ccgcatacac tattctcaga atgacttggg tgagtactca ccagtcacag 2940  
aaaagcatct tacggatggc atgacagtaa gagaattatg cagtgtgcc ataaccatga 3000  
gtgataacac tgcggccaac ttacttctga caacgatcgg aggaccgaag gagctaaccg 3060  
cttttttgca caacatgggg gatcatgtaa ctgccttga tcgttgggaa ccggagctga 3120  
atgaagccat accaaacgac gagcgtgaca ccacgatgcc tgtagcaatg gcaacaacgt 3180  
tgcgcaaact attaaactggc gaactactta ctctagcttc ccggcaacaa ttaatagact 3240  
ggatggaggc ggataaagtt gcaggaccac ttctgcgctc ggcccttccg gctggctggg 3300  
ttattgctga taaatctgga gccggtgagc gtgggtctcg cggatcatt gcagcactgg 3360  
ggccagatgg taagccctcc cgtatcgtag ttatctacac gacggggagt caggcaacta 3420  
tggtgaacg aaatagacag atcgttgaga taggtgcctc actgattaag cattggtaac 3480  
tgtcagacca agtttactca tatatacttt agattgattt aaaacttcat ttttaattta 3540  
aaaggatcta ggtgaagatc ctttttgata atctcatgac caaaatccct taacgtgagt 3600  
tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa aggatcttct tgagatcctt 3660  
ttttctgcg cgtaactctgc tgcttgcaaa caaaaaaacc accgctacca gcggtgggtt 3720  
gtttgccgga tcaagagcta ccaactcttt ttccgaaggg aactggcttc agcagagcgc 3780  
agataccaaa tactgtcctt ctagtgtagc cgtagttagg ccaccacttc aagaactctg 3840  
tagcaccgcc tacatacctc gctctgctaa tctgttacc agtggctgct gccagtggcg 3900

37

```

ataagtcgtg tcttaccggg ttggactcaa gacgatagtt accggataag ggcgagcggg 3960
cgggctgaac ggggggttcg tgcacacagc ccagcttgga gcgaacgacc tacaccgaac 4020
tgagatacct acagcgtgag ctatgagaaa gcgccacgct tcccgaaggg agaaaggcgg 4080
acaggatatc ggtaagcggc agggctcggaa caggagagcg cacgagggag cttccagggg 4140
gaaacgcctg gtatctttat agtcctgtcg ggtttcgcca cctctgactt gagegtcgat 4200
ttttgtgatg ctcgtcaggg gggcggagcc tatggaaaaa cgccagcaac ggggcctttt 4260
tacggttcct ggccttttgc tggccttttg ctcacatggt ctttctcgcg ttatccctcg 4320
attcattaat gcaggtcacg atc                                     4343

```

&lt;210&gt; 9

&lt;211&gt; 3888

&lt;212&gt; DNA

&lt;213&gt; Influenza virus

&lt;400&gt; 9

```

ccccaaaaaa aaaaaaaaaa aaaaaaaaaa agtccagagt ggccccgcgg ttccgcgcgg 60
gggggggggg ggggggggga cactttcggg catctggtcg acctccagca tggggggaaa 120
aaaaaaaaac aaagtttcgc ccggagtact ggtcgacctc cgaagtggg ggggagtaga 180
aacagggtag ataatcactc actgagtgc atccacatcg cgagcgcgcg taatacgact 240
cactataggg cgaattgggt accgggcccc cctcagaggt cgacggtatc gataagcttc 300
gacgagattt tcaggagcta aggaagctaa aatggagaaa aaaatcactg gatataccac 360
cgttgatata tcccaatggc atcgtaaaga acattttgag gcatttcagt cagttgctca 420
atgtacctat aaccagaccg ttcagctgga tattacggcc tttttaaaga ccgtaaagaa 480
aaataagcac aagttttatc cggcctttat tcacattctt gccgcctga tgaatgctca 540
tccggaattc cgtatggcaa tgaaagacgg tgagctgggtg atatgggata gtgttcaccc 600
ttgttacacc gttttccatg agcaaaactga aacgttttca tcgctctgga gtgaatacca 660
cgacgatttc cggcagtttc tacacatata ttcgcaagat gtggcgtgtt acggtgaaaa 720
cctggcctat ttccctaaag ggtttattga gaatatgttt ttcgtctcag ccaatccctg 780
ggtgagtttc accagttttg atttaaacgt ggccaatatg gacaacttct tcgccccctg 840
tttcaccatg ggcaaatatt atacgaagg cgacaagggtg ctgatgccgc tggcgattca 900
ggttcacatc gccgtttgtg atggcttcca tgcggcaga atgcttaatg aattacaaca 960
gtactgcgat gagtggcagg gcggggcgta atttttttaa ggcagttatt ggtgccctta 1020
aacgcctggt gctacgctg aataagtgat aataagcgga tgaatggcag aaattcgtcg 1080
aagcttgata tcgaattcct gcagcccggg ggatccacta gttctagagc ggccgccacc 1140
gcggtggagc tccagctttt gttcccttta gtgaggggta attgcgcgca ggcctagcta 1200
ggtaaagaaa aatacccttg attcttctaa taaccggcg gcccaaatg ccgactcgga 1260
gcgaaagata tacctcccc ggggcccggg ggtcgcgtca ccgaccacgc cgccggccca 1320
ggcgacgcgc gacacggaca cctgtcccca aaaacgccac catcgagcc acacacggag 1380
cgcccggggc cctctggtca accccaggac acacgcggga gcagcgcggg gccggggacg 1440
ccctcccggc cgcccggtgc acacgcaggg ggccggcccc tgtctccaga gcgggagccg 1500
gaagcatttt cgccggcccc ctctacgac cgggacacac gagggaccga aggccggcca 1560
ggcgcgacct ctcgggccgc acgcgcgctc agggagcgtc ctccgactcc gcacggggac 1620
tcgccagaaa ggatcgtgac ctgcattaat gaatcagggg ataacgcagg aaagaacatg 1680
tgagcaaaaag gccagcaaaa ggccagggaac cgtaaaaagg ccgcgttgct ggcgtttttc 1740
cataggctcc gccccctga cgagcatcac aaaaatcgac gctcaagtca gaggtggcga 1800
aaccgcagag gactataaag ataccaggcg tttccctctg gaagctccct cgtgcgtctc 1860
cctgttccga ccctgccgt taccggatac ctgtccgctt ttctcccttc gggaagcgtg 1920
gcgctttctc atagctcacg ctgtaggtat ctcagttcgg tgtaggtcgt tcgctccaag 1980
ctgggctgtg tgcacgaacc ccccggttcag cccgaccgct gcgccttatc cggtaactat 2040
cgtcttgagt ccaaccgggt aagacacgac ttatcgccac tggcagcagc cactggtaac 2100
aggattagca gagcgaggta tgtaggcggg gctacagagt tcttgaagtg gtggcctaac 2160
tacggctaca ctagaaggac agtatttggg atctgcgctc tgcgtgaagc agttaccttc 2220
ggaaaaagag ttggtagctc ttgatccggc aaacaaacca ccgctggtag cgggtggttt 2280
tttgtttgca agcagcagat tacgcgcaga aaaaaggat ctcaagaaga tcctttgatc 2340
ttttctacgg ggtctgacgc tcagtggaaac gaaaactcac gttaagggat tttggtcatg 2400

```

38

agattatcaa	aaaggatctt	cacctagatc	cttttaaatt	aaaaatgaag	ttttaaatca	2460
atctaaagta	tatatgagta	aacttgggtc	gacagttacc	aatgcttaat	cagtgaaggca	2520
cctatctcag	cgatctgtct	atttcgttca	tccatagttg	cctgactccc	cgctcgtgtag	2580
ataactacga	tacgggaggg	cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac	2640
ccacgctcac	cggtccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	2700
agaagtggtc	ctgcaacttt	atccgctccc	atccagtcta	ttaattgttg	ccgggaagct	2760
agagtaagta	gttcgccagt	taatagtttg	cgcaacgttg	ttgccattgc	tacaggcatc	2820
gtggtgtcac	gctcgtcgtt	tggatgggt	tcattcagct	ccggttccca	acgatcaagg	2880
cgagttacat	gatccccat	gttgtgcaaa	aaagcggtta	gtccttcggg	tcctccgata	2940
gttgtcagaa	gtaagtggc	cgcagtgtta	tcactcatgg	ttatggcagc	actgcataat	3000
tctcttactg	tcatgccatc	cgtaagatgc	ttttctgtga	ctggtgagta	ctcaaccaag	3060
tcattctgag	aatagtgtat	gcggcgaccg	agttgctctt	gcccggcgct	aacacgggat	3120
aataccgcgc	cacatagcag	aactttaaaa	gtgttcataa	ttggaaaacg	ttcttcgggg	3180
cgaaaactct	caaggatctt	accgctgttg	agatccagtt	cgatgtaacc	cactcgtgca	3240
cccaactgat	cttcagcatc	ttttactttc	accagcgttt	ctgggtgagc	aaaaacagga	3300
aggcaaaatg	ccgcaaaaaa	gggaataagg	gcgacacgga	aatgttgaat	actcatactc	3360
ttcctttttc	aatattattg	aagcatttat	cagggttatt	gtctcatgag	cggatacata	3420
tttgaatgta	tttagaaaaa	taaacaaaag	agtttgtaga	aacgcaaaaa	ggccatccgt	3480
caggatggcc	ttctgcttaa	tttgatgcct	ggcagtttat	ggcgggcgct	ctgcccgcga	3540
ccctccgggc	cgttgcttcg	caacgttcaa	atccgctccc	ggcggatttg	tcctactcag	3600
gagagcggtc	accgacaaac	aacagataaa	acgaaaggcc	cagtctttcg	actgagcctt	3660
tcgtttttat	tgatgcctgg	cagttcccta	ctctcgcata	gggagacccc	acactaccat	3720
cggcgctacg	gcgtttcact	tctgagttcg	gcattggggc	aggtggggacc	accgcgctac	3780
tgccgccagg	caaattctgt	tttatcagac	cgcttctgcg	ttctgattta	atctgtatca	3840
ggctgaaaat	cttctctcat	ccgccaaaac	agaagctagc	ggccgatac		3888



PO - Munich  
33  
06. März 1999

## Claims

1. A recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus; wherein  
(a) at least one of the regular viral RNA segments has been exchanged for a vRNA encoding a foreign gene; and/or  
~~(b) at least one of the regular viral RNA segments is an ambisense RNA molecule, containing~~  
one of the standard viral genes in sense orientation and a foreign, recombinant gene in anti-sense orientation, or *vice versa*, in overall convergent arrangement.
2. The recombinant virus according to claim 1, wherein in the ambisense RNA molecule said foreign recombinant gene is covalently bound to one of the viral genes, while the original vRNA segment coding for the same gene is deleted from the recombinant virus by way of specific ribozyme cleavage.
3. The recombinant virus according to claims 1 and 2 in which one or both of the standard glycoproteins hemagglutinin and neuraminidase have been exchanged into foreign glycoprotein(s) or into fusion glycoproteins consisting of an anchor segment derived from hemagglutinin and an ectodomain obtained from the foreign source, viral or cellular, or in which such recombinant glycoprotein has been inserted as a third molecular species in addition to the remaining standard components.
4. The recombinant virus according to claims 1 to 3, in which the terminal viral RNA sequences, which are active as the promoter signal, have been varied by nucleotide substitutions in up to five positions, resulting in improved transcription rates of both the vRNA promoter as well as the cRNA promoter as present in the complementary sequence.
5. The recombinant virus according to claims 1 to 4, in which the foreign gene(s) in ambisense covalent junction with viral gene(s) code for proteins and/or glycoproteins which are secreted from cells infected with the recombinant virus.

6. The recombinant virus according to claims 1 to 4, in which the foreign gene(s) in ambisense covalent junction with viral gene(s) code for proteins or artificial polypeptides designed to support an efficient presentation of inherent epitopes at the surface of (abortively) infected cells, for stimulation of a B cell and /or T cell response.
7. A method for the production of recombinant influenza viruses as defined in claims 1 to 6 comprising
- 
- (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, in antisense, sense or ambisense design,
- (b) followed by infection with an influenza carrier strain constructed to include flanking ribozyme target sequences in at least one of its viral RNA segments, and
- (c) thereafter selective vRNA inactivation through ribozyme cleavage.
8. A method of constructing influenza carrier strains carrying one or more ribozyme target sites (of type one) in vRNA flanking positions comprising
- (a) RNA polymerase I synthesis of recombinant vRNAs *in vivo*, carrying two different 3' promoter sequences in tandem, which are separated by a second type of ribozyme target sequence, and which carry the said internal ribozyme target sites of type one;
- (b) followed by infection of an influenza wildtype strain;
- (c) thereafter amplification through simple steps of viral propagation; and
- (d) finally isolation through removal of their external 3' promoter sequence by ribozyme cleavage through infection of cells expressing ribozyme type 2, followed by plaque purification.
9. A ribozyme-sensitive influenza carrier strain obtainable by the method of claim 8.
10. A pharmaceutical composition comprising a recombinant virus according to claims 1 to 6.
11. Use of a recombinant virus according to claims 1 to 6 for preparing a medicament for vaccination purposes.

12. The use according to claim 11, wherein the medicament
- (a) is suitable against influenza and/or against other infections;
  - (b) is present in form of inactivated preparations; and/or
  - (c) is present in form of live recombinant (attenuated) viruses.

13. Use of a recombinant virus according to claims 1 to 6 as vector systems in somatic gene therapy, ~~for transfer and expression of foreign genes into cells (abortively) infected by such~~ viruses, either in *ex vivo* or *in vivo* application schemes.

14. A method for the production of proteins or glycoproteins which comprises utilizing a recombinant influenza virus according to claims 1 to 6 as expression vector.

15. The method of claim 14, wherein the production is performed in cell culture cells or in fertilized chicken eggs.

EPO - Munich  
33  
06. März 1999

FIG. 1




FIG. 2





FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7






FIG. 8




FIG. 9

**cRNA-promoter**









FIG. 11

A



bp-variant position	G - C	A - U	C - G	U - A
2 - 9	pHL.2024 100%	pHL.1921 41%	pHL.2003 < 3%	pHL.2004 < 3%
3 - 8	pHL.2002 30%	pHL.1920 121%	pHL.1148 39%	pHL.2024 100%
2 - 9	pHL.1945 11%	pHL.1946 30%	pHL.2024 100%	pHL.1923 28%
3 - 8	pHL.2428 6%	pHL.2024 100%	pHL.1948 33%	pHL.1922 97%

B



C




FIG. 12




FIG. 13




FIG. 14






FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19





FIG. 20



EPO - Munich  
33  
06. März 1999Abstract

## Stable Recombinant Influenza Viruses Free of Helper Viruses

The invention relates to a recombinant influenza virus for high-yield expression of incorporated foreign gene(s), which is genetically stable in the absence of any helper virus; wherein

~~(a) at least one of the regular viral RNA segments has been exchanged for a vRNA encoding a foreign gene; and/or~~

(b) at least one of the regular viral RNA segments is an ambisense RNA molecule, containing one of the standard viral genes in sense orientation and a foreign, recombinant gene in anti-sense orientation, or *vice versa*, in overall convergent arrangement.

The present invention further provides a method for the production of said recombinant influenza virus, a method for constructing specific ribozyme-sensitive influenza carrier strains; pharmaceutical compositions comprising said recombinant influenza virus; and the use of said recombinant influenza virus for preparing medicaments for vaccination purposes.